


Volume 141 No. 1 May 2006 ISSN 0269-7491



ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning


ASSOCIATE EDITORS

K.E. Havens

S.V. Krupa

K.C. Jones

J.W. Erisman



ENVIRONMENTAL POLLUTION

CONTENTS—Continued from outside back cover

- 98 **Comparative kinetic desorption of ^{60}Co , ^{85}Sr and ^{134}Cs from a contaminated natural silica sand column: Influence of varying physicochemical conditions and dissolved organic matter**
N. Solovitch-Vella, J.-M. Garnier
Desorption of RN(s) was related to chemical characteristics of organic colloids.
- 107 **Distributions and Concentrations of PAHs in Hong Kong Soils**
H.B. Zhang, Y.M. Luo, M.H. Wong, Q.G. Zhao, G.L. Zhang
Baseline information is provided on levels, distributions and possible sources of PAHs in Hong Kong soils.
- 115 **Field evaluation of Cd and Zn phytoextraction potential by the hyperaccumulators *Thlaspi caerulescens* and *Arabidopsis halleri***
S.P. McGrath, E. Lombi, C.W. Gray, N. Caille, S.J. Dunham, F.J. Zhao
*The hyperaccumulator *Thlaspi caerulescens* (the Ganges ecotype) is more efficient at phytoextracting Cd than Zn from contaminated soil.*
- 126 **A link between lead and cadmium kinetic speciation in seawater and accumulation by the green alga *Ulva lactuca***
J.O. Muse, C.N. Carducci, J.D. Stripeikis, M.B. Tudino, F.M. Fernández
Metal kinetic speciation can overcome CF limitations for the evaluation of marine pollution.
- 131 **Effects of nitrogen with and without acidified sulphur on an ectomycorrhizal community in a Sitka spruce (*Picea sitchensis* Bong. Carr) forest**
J.A. Carfrae, K.R. Skene, L.J. Sheppard, K. Ingleby, A. Crossley
*N deposition effects on *T. fibrillosa* versus *L. rufus* colonisation of a Sitka spruce plantation.*
- 139 **Simulating the long-term chemistry of an upland UK catchment: Heavy metals**
E. Tipping, A.J. Lawlor, S. Lofts, L. Shotbolt
Catchment turnover times for the strongly-retained metals Cu and Pb are of the order of centuries, whereas those for the more mobile Ni, Zn and Cd are appreciably shorter.
- 151 **Simulating the long-term chemistry of an upland UK catchment: Major solutes and acidification**
E. Tipping, A.J. Lawlor, S. Lofts
CHUM-AM accounts for recovery from acidification due to sulphur deposition, but predicts re-acidification if nitrogen deposition is not reduced.
- 167 **Responses to ammonium and nitrate additions by boreal plants and their natural enemies**
A. Nordin, J. Strengbom, L. Ericson
Biota will respond to nitrogen deposition depending on the form of nitrogen.
- 175 **Seasonal development of ozone-induced foliar injury on tall milkweed (*Asclepias exaltata*) in Great Smoky Mountains National Park**
L. Souza, H.S. Neufeld, A.H. Chappelka, K.O. Burkey, A.W. Davison
Leaf loss was not necessarily accompanied by symptoms of foliar ozone injury.
- 184 **Sorption, desorption and mineralisation of the herbicides glyphosate and MCPA in samples from two Danish soil and subsurface profiles**
S.R. Sørensen, A. Schultz, O.S. Jacobsen, J. Aamand
Natural attenuation potential of the herbicides glyphosate and MCPA was assessed in soil and subsurface profiles.

CONTENTS

Volume 141 Number 1 2006

- 1 Acute toxicity of heavy metals towards freshwater ciliated protists**
P. Madoni, M.G. Romeo
Ciliated protozoa are suitable bioindicators of heavy metal pollution in freshwater environments.
- 8 Assessment of ozone effects on nitrate export from Hubbard Brook Watershed 6**
B. Hong, D.A. Weinstein, D.P. Swaney
Ambient ozone was estimated to cause an additional 0.042 gN/m² per year of nitrate to be exported from Hubbard Brook Watershed 6 during 1964–1994.
- 22 Extraction of arsenate and arsenite species from soils and sediments**
M. Georgiadis, Y. Cai, H.M. Solo-Gabriele
NaDDC preserves arsenite during soil and sediment extraction.
- 30 Organohalogen chemicals in human blood from the United Kingdom**
G.O. Thomas, M. Wilkinson, S. Hodson, K.C. Jones
A survey of PBDE, PCB and OC pesticide concentrations in human blood helps identify baseline concentrations in the UK population and found decabromodiphenyl ether in non-occupationally exposed individuals.
- 42 Response of drinking-water reservoir ecosystems to decreased acidic atmospheric deposition in SE Germany: Trends of chemical reversal**
K.-U. Ulrich, L. Paul, A. Meybohm
Trend analysis demonstrates significant chemical response of acidified surface waters to intensely decreased acidic atmospheric deposition from formerly high levels in the Erzgebirge (SE Germany).
- 54 Use of carbon and nitrogen stable isotope ratios to assess the effects of environmental contaminants on aquatic food webs**
T. Caquet
Carbon and nitrogen stable isotope ratios may be used to detect the impact of pollutants on aquatic food webs.
- 60 Evidence of endocrine alteration in the red mullet, *Mullus barbatus* from the NW Mediterranean**
R. Martin-Skilton, R. Lavado, R. Thibaut, C. Minier, C. Porte
Red mullet may be more susceptible to endocrine disruptors during the reproductive period.
- 69 Lead and zinc accumulation and tolerance in populations of six wetland plants**
H. Deng, Z.H. Ye, M.H. Wong
Populations from metal contaminated sites did not have significantly higher metal tolerance indices.
- 81 Recovery of the histogram of hourly ozone distribution from weekly average concentrations**
L.E. Olcese, B.M. Toselli
The histogram of hourly ozone distribution can be obtained based on passive sensor data.
- 90 Interaction of chemical cues from fish tissues and organophosphorous pesticides on *Ceriodaphnia dubia* survival**
J.D. Maul, J.L. Farris, M.J. Lydy
*Potentiation of organophosphorous pesticide toxicity to *Ceriodaphnia dubia* by fathead minnow (*Pimephales promelas*) chemical cues was observed.*