

Volume 141 No. 3 June 2006 ISSN 0269-7491

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning


ASSOCIATE EDITORS

K.E. Havens

S.V. Krupa

K.C. Jones

J.W. Erisman



ENVIRONMENTAL POLLUTION

CONTENTS—Continued from outside back cover

- 469 Sequential extraction combined with isotope analysis as a tool for the investigation of lead mobilisation in soils: Application to organic-rich soils in an upland catchment in Scotland**
J.R. Bacon, J.G. Farmer, S.M. Dunn, M.C. Graham, S.I. Vinogradoff
Sequential extraction combined with isotope analysis was used as a tool to assess mobilisation of lead into streams.
- 482 Effects of oxygen on the release and distribution of phosphorus in the sediments under the light condition**
X. Jiang, X. Jin, Y. Yao, L. Li, F. Wu
Uptake of dissolved inorganic P affected P release and distribution.
- 488 Factors influencing arsenic accumulation by *Pteris vittata*: A comparative field study at two sites**
C.Y. Wei, X. Sun, C. Wang, W.Y. Wang
*Nutrient elements influenced the arsenic-accumulation capacity of *Pteris vittata* in the field.*
- 494 Linking organic pollutant (bio)availability with geosorbent properties and biomimetic methodology: A review of geosorbent characterisation and (bio)availability prediction**
G.A.C. Ehlers, A.P. Loibner
Hydrophobic organic pollutant retention in geosorbent matrices, soil and sediment characteristics, biomimetic techniques is reviewed.
- 513 Mercury in fish from two Nicaraguan lakes: A recommendation for increased monitoring of fish for international commerce**
J.K. McCrary, M. Castro, K.R. McKaye
Fish commonly exported into the international food supply may be contaminated with mercury.
- 519 Effects of polycyclic aromatic hydrocarbons on germination and subsequent growth of grasses and legumes in freshly contaminated soil and soil with aged PAHs residues**
M.J. Smith, T.H. Flowers, H.J. Duncan, J. Alder
Germination tests alone do not predict growth of plants on PAH contaminated soil.
- 526 Effects of added PAHs and precipitated humic acid coatings on phenanthrene sorption to environmental Black carbon**
G. Cornelissen, Ö. Gustafsson
Sorption of phenanthrene to Black carbon is attenuated by other PAHs but not by humic acid coatings.
- 532 Low levels of persistent organic pollutants (POPs) in New Zealand eels reflect isolation from atmospheric sources**
N. Holmqvist, P. Stenroth, O. Berglund, P. Nyström, K. Olsson, D. Jellyman, A.R. McIntosh, P. Larsson
Low levels of PCBs found in New Zealand eels reflect isolation from atmospheric sources while DDTs levels are determined by land use.
- 539 Efficient retrieval of vegetation leaf area index and canopy clumping factor from satellite data to support pollutant deposition assessments**
N. Nikolov, K. Zeller
The paper presents a physics-based algorithm for retrieval of vegetation LAI and canopy-clumping factor from satellite data to assist research of pollutant deposition and trace-gas exchange. The method is employed to derive a monthly LAI dataset for the conterminous USA and verified at a continental scale.
- 550 Export of nutrients from golf courses on the Precambrian Shield**
J.G. Winter, P.J. Dillon
Golf courses increase nutrient loads in receiving streams.
- 555 The status of pesticide pollution in surface waters (rivers and lakes) of Greece. Part I. Review on occurrence and levels**
I.K. Konstantinou, D.G. Hela, T.A. Albanis
Information on pesticide pollution of surface waters in Greece is reviewed.

CONTENTS

Volume 141 Number 3 2006

Invited papers

- 387 Woody biomass phytoremediation of contaminated brownfield land**
C.J. French, N.M. Dickinson, P.D. Putwain
Field trials show short-rotation coppice provides effective risk management and remediation solutions to hotspots of residual metal and As contamination of brownfield land.
- 396 Eisenia fetida increased removal of polycyclic aromatic hydrocarbons from soil**
S.M. Contreras-Ramos, D. Álvarez-Bernal, L. Dendooven
Addition of earthworms, Eisenia fetida, accelera removal of polycyclic aromatic hydrocarbons from soil.
- 402 Identifying primary stressors impacting macroinvertebrates in the Salinas River (California, USA): Relative effects of pesticides and suspended particles**
B.S. Anderson, B.M. Phillips, J.W. Hunt, V. Connor, N. Richard, R.S. Tjeerdema
Pesticides are the primary stressor impacting macroinvertebrates in sections of the lower Salinas River.
- 409 Growth of the damselfly Ischnura heterosticta is better in saline water than freshwater**
B.J. Kefford, L. Zaluzniak, D. Nugegoda
This species while typically found in freshwaters does better in saline waters.

Regular papers

- 420 Effects of phosphorus fertilization on the availability and uptake of uranium and nutrients by plants grown on soil derived from uranium mining debris**
G. Rufyikiri, J. Wannijn, L. Wang, Y. Thiry
Phosphate addition to soil derived from uranium mining debris may decrease U mobility and bioavailability to plant.
- 428 An exploration of the relationship between adsorption and bioavailability of pesticides in soil to earthworm**
Y.L. Yu, X.M. Wu, S.N. Li, H. Fang, H.Y. Zhan, J.Q. Yu
Bioavailability of pesticides in soil to earthworm is governed by adsorption characteristics.
- 434 Field assessment of oxytetracycline exposure to the freshwater macrophytes Egeria densa Planch. and Ceratophyllum demersum L.**
M.L. Hanson, C.W. Knapp, D.W. Graham
Oxytetracycline did not pose a risk for two freshwater macrophytes.
- 443 Responses of antioxidant systems and LPO level to benzo(a)pyrene and benzo(k)fluoranthene in the haemolymph of the scallop Chlamys ferrari**
L.Q. Pan, J. Ren, J. Liu
Biomarkers for PAHs were identified in the scallop Chlamys ferrari.
- 452 Response of the Mediterranean sponge Chondrosia reniformis Nardo to copper pollution**
E. Cebrian, G. Agell, R. Martí, M.J. Uriz
Contrasting effects of copper pollution on the Atlanto-Mediterranean sponge Chondrosia reniformis.
- 459 Degradation of polycyclic aromatic hydrocarbons (PAHs) in an aged coal tar contaminated soil under in-vessel composting conditions**
B. Antizar-Ladislao, J. Lopez-Real, A.J. Beck
Maximum degradation of PAHs in an aged coal tar contaminated soil can be achieved using optimal operational conditions during composting.