

530 AGRO


530

The contents of this document are copyright works and unless you have the permission of copyright owner or of The Copyright Licensing Agency Ltd or another authorised licensing body or except as may be permitted by statute this document may not be copied (including storage in any electronic medium) or otherwise reproduced (even for internal purposes) or resold.

LIBRARY  
LIBRARY  
Supplied by, or on behalf of  
THE BRITISH LIBRARY  
Document Supply Service  
Boston Spa  
Wetherby  
West Yorkshire  
LS23 7BQ  
United Kingdom

N° 31/03.

Agro.



#### LITERATURE CITED

1. Alström, S. 1991. Induction of disease resistance in common bean susceptible to halo blight bacterial pathogen after seed bacterization with rhizosphere pseudomonads. *J. Gen. Appl. Microbiol.* 37:495-501.
2. Brian, P. W. 1957. The ecological significance of antibiotic production. Pages 168-188 in: *Microbiological Ecology*. R. E. O. Williams and C. C. Spencer, eds. Cambridge University Press, Cambridge.
3. Caruso, F. L., and Kuć, J. 1977. Field protection of cucumber, watermelon, and muskmelon against *Colletotrichum lagenarium* by *Colletotrichum lagenarium*. *Phytopathology* 67:1290-1292.
4. Cox, J. E., Fraser, L. R., and Broadbent, P. 1976. Stem pitting of grape fruit. Field protection by the use of mild strains: An evaluation of trials in two climatic districts. Pages 68-70 in: *Proc. Conf. Int. Organ. Citrus Virol.*, 7th. E. C. Calvin, ed. University of Florida Press, Gainesville.
5. Daly, J. M. 1972. The use of near-isogenic lines in biochemical studies of the resistance of wheat to stem rust. *Phytopathology* 62:392-400.
6. Deacon, J. W. 1991. Significance of ecology in the development of biocontrol agents against soil-borne plant pathogens. *Biocontrol Sci. Technol.* 1:5-20.
7. Dean, R. A., and Kuć, J. 1985. Induced systemic protection in plants. *Trends Biotechnol.* 3:125-129.
8. Dehne, H. W., Stenzel, K., and Schonbeck, F. 1984. The efficiency of induced resistance under practical culture conditions. III. Reproduction of powdery mildew on induced resistant plants. *J. Plant Dis. Prot.* 91:258-265.
9. Doke, N., Garas, N. A., and Kuć, J. 1979. Partial characterization and aspects of the mode of action of a hypersensitivity-inhibiting factor (HIF) isolated from *Phytophthora infestans*. *Physiol. Plant Pathol.* 15:127-140.
10. Garas, N. A., Doke, N., and Kuć, J. 1979. Suppression of the hypersensitive reaction in potato tubers by mycelial components from *Phytophthora infestans*. *Physiol. Plant Pathol.* 15:117-126.
11. Hammerschmidt, R., and Kuć, J. 1982. Lignification as a mechanism for induced systemic resistance in cucumber. *Physiol. Plant Pathol.* 20:61-71.
12. Hammerschmidt, R., Lamport, D. T. A., and Muldoon, E. P. 1984. Cell wall hydroxyproline enhancement and lignin deposition as an early event in the resistance of cucumber to *Cladosporium cucumerinum*. *Physiol. Plant Pathol.* 24:43-47.
13. Hammerschmidt, R., Nuckles, E., and Kuć, J. 1982. Association of enhanced peroxidase activity with induced systemic resistance of cucumber to *Colletotrichum lagenarium*. *Physiol. Plant Pathol.* 20:73-82.
14. Heath, M. C. 1982. The absence of active defense mechanisms in compatible host-pathogen interactions. Pages 143-156 in: *Active Defense Mechanisms in Plants*. R. K. S. Wood, ed. Plenum Press, New York.
15. Kloepper, J. W., Tuzun, S., and Kuć, J. 1992. Proposed definitions related to induced disease resistance. *Biocontrol Sci. Technol.* 2:349-351.
16. Kuć, J. 1983. Induced systemic resistance in plants to diseases caused by fungi and bacteria. Pages 191-221 in: *The Dynamics of Host Defense*. J. A. Bailey and B. J. Deverall, eds. Academic Press, Ltd., Sydney, Australia.
17. Kuć, J., and Preisig, C. 1984. Fungal regulation of disease resistance mechanisms in plants. *Mycologia* 76:767-784.
18. Kuć, J., and Rush, J. S. 1985. Phytoalexins. *Arch. Biochem. Biophys.* 236:455-472.
19. Lawton, M. A., and Lamb, C. J. 1987. Transcriptional activation of plant defense genes by fungal elicitor, wounding and infection. *Mol. Cell. Biol.* 7:335-341.
20. Liu, L., Kloepper, J. W., and Tuzun, S. 1995. Induction of systemic resistance in cucumber against Fusarium wilt by plant growth-promoting rhizobacteria. *Phytopathology* 85:695-698.
21. Liu, L., Kloepper, J. W., and Tuzun, S. 1995. Induction of systemic resistance in cucumber against bacterial angular leaf spot by plant growth-promoting rhizobacteria. *Phytopathology* 85:843-847.
22. Rothrock, C. S., and Gottlieb, D. 1984. Role of antibiosis in antagonism of *Streptomyces hygroscopicus* var. *geldanus* to *Rhizoctonia solani* in soil. *Can. J. Microbiol.* 30:1440-1447.
23. Sasser, M. 1990. Identification of bacteria through fatty acid analysis. Pages 199-204 in: *Methods in Phytobacteriology*. Z. Klement, K. Rudolph, and D. C. Sands, eds. Akademiai Kiad, Budapest.
24. Tuzun, S., Juarez, J., Nesmith, W. C., and Kuć, J. 1992. Induction of systemic resistance in tobacco against metalaxyl-tolerant strains of *Peronospora tabacina* and the natural occurrence of the phenomenon in Mexico. *Phytopathology* 82:425-429.
25. Tuzun, S., and Kuć, J. 1991. Plant immunization: An alternative to pesticides for control of plant diseases in the greenhouse and field. Pages 30-40 in: *The Biological Control of Plant Diseases - Proceedings of the International Seminar "Biological Control of Plant Diseases and Virus Vectors"*. J. Bay-Petersen, ed. Food and Fertilizer Technology Center, Taipei, Republic of China.
26. Tuzun, S., Nesmith, W., Ferriss, R. S., and Kuć, J. 1986. Effects of stem injections with *Peronospora tabacina* on growth of tobacco and protection against blue mold in the field. *Phytopathology* 76:938-941.
27. Van Peer, R., Niemann, G. J., and Schippers, B. 1991. Induced resistance and phytoalexin accumulation in biological control of Fusarium wilt of carnation by *Pseudomonas* sp. strain WCS417r. *Phytopathology* 81:728-734.
28. Wei, G., Kloepper, J. W., and Tuzun, S. 1991. Induction of systemic resistance of cucumber to *Colletotrichum orbiculare* by select strains of plant growth-promoting rhizobacteria. *Phytopathology* 81:1508-1512.
29. Zhou, T., and Paulitz, T. C. 1994. Induced resistance in the biocontrol of *Pythium aphanidermatum* by *Pseudomonas* spp. on cucumber. *J. Phytopathol.* 142:51-63.



## Correlation Between Susceptibility to Crown Gall and Sensitivity to Cytokinin in Aspen Cultivars

T. Beneddra, C. Picard, A. Petit, and X. Nesme

First author: Station de Pathologie Végétale et Phytobactériologie, INRA, Centre de Recherches d'Angers, 49000 Beaucouzé, France; second and fourth authors: Laboratoire de Microbiologie des Sols, URA CNRS 1977, and INRA, Université Lyon 1, 69622 Villeurbanne Cedex, France; and third author: Institut des Sciences Végétales, CNRS, 91198 Gif-sur-Yvette, France.

This study was supported by the Forest Contract MA1B006C from the European Union for X. Nesme, and by a scholarship from INRA for C. Picard. We thank J. P. Hernalsteens and A. Reynaerts for supplying T-DNA mutants, and D. Cornu for aspen shoot cultures.

Accepted for publication 2 November 1995.

### ABSTRACT

Beneddra, T., Picard, C., Petit, A., and Nesme, X. 1996. Correlation between susceptibility to crown gall and sensitivity to cytokinin in aspen cultivars. *Phytopathology* 86:225-231.

Closely related aspen cultivars (*Populus tremula* × *P. alba*) were ranked as resistant, intermediate, or susceptible to crown gall according to their tumor response to the selected strains C58, B6, and 354 of *Agrobacterium tumefaciens*, respectively. As shown with mutant agrobacteria that harbored derivatives of pTiB6S3, a strain retained its ability to define differences in susceptibility among aspen cultivars as long as it had a functional *ipt* gene and, therefore, could induce the biosynthesis of a

cytokinin in transformed plants. This suggested that differences in susceptibility to crown gall were related to differences in sensitivity to cytokinin. Cultivar sensitivity to cytokinin was determined in vitro by a leaf disk assay. Aspen cultivars resistant to the highest cytokinin concentrations (32 μM benzyladenine) were also found to be resistant to *A. tumefaciens* strain B6 and the most resistant to natural crown gall infections. This result showed that sensitivity to cytokinin was a plant factor controlling tumorigenesis.

*Additional keywords:* hormone, host range, *iaa*, poplar, T-DNA mutants.

The crown gall disease is characterized by tissue overgrowths generally located on the crown and roots of dicots, gymnosperms, and some monocots (11). It is a major disease that is widespread in tree nurseries throughout the world, affects growth and commercial quality of stocks of many species (2), and causes important financial losses in orchard production (14,30). In some instances, breeders have abandoned species or cultivars that are too susceptible to crown gall (27). Such a decision is particularly serious when the plants are required for special purposes, such as aspen hybrids (*Populus tremula* × *P. alba*) that have been selected for forestation of hydromorphic soils (21). Crown gall is common in aspen nurseries (31), and various means of control have been attempted, including biocontrol with strain K84 and culling of diseased plants. However, biological control had no or very limited effects, probably because K84-resistant agrobacteria were already present in the nursery soil (X. Nesme, unpublished data). Culling was more efficient, but did not eradicate the disease (30). Thus, the selection of resistant aspens has also been considered.

Although resistant genotypes have been selected in several host species (3,26,27,36,47,54), plant traits that control tumorigenesis are poorly understood compared with present knowledge of virulence in *Agrobacterium* spp. (37,51). Most bacterial tumorigenic functions are encoded by a tumor-inducing plasmid or Ti plasmid (50). During the infection process, a fragment of this plasmid, called T-DNA, is transferred to the plant genome, where it is expressed like native plant genes (10). Transfer of T-DNA is controlled by virulence genes located in the nontransferred *vir* region of the Ti plasmid upon induction of compounds exuded by wounded plant cells (45). Oncogenes responsible for tumor

growth reside in the T-DNA region and cause overproduction of auxin (*iaaM* and *iaaH*) and cytokinin (*ipt*) in transformed cells (1,17,42). Bacterial host range is mainly controlled by the Ti plasmid (19,23,48). Studies have shown that the *vir* genes or the T-DNA oncogenes, particularly the *ipt* gene, may modify the host range to some extent (5,16,52,53). Differences in resistance between hosts can, therefore, be caused by plant factors that interact with *vir* genes or T-DNA oncogenes (12,29,46).

In this paper, a large number of wild-type strains of *A. tumefaciens* were screened for their ability to define differences in susceptibility among closely related cultivars of *P. tremula* × *P. alba*. Research with mutant *A. tumefaciens* was undertaken to show whether *iaa* and *ipt* genes were required to define cultivar differences. Results indicated a role for *ipt*, that encodes the biosynthesis of a cytokinin in transformed plants. Thus, the sensitivity of host tissues to exogenous cytokinin was estimated by a leaf disk assay, and sensitivity to cytokinin was compared with crown gall susceptibility after artificial and natural infections.

### MATERIALS AND METHODS

**Bacteria.** A large number of wild-type strains of *Agrobacterium* spp., provided by the Collection Française de Bactéries Phytopathogènes (CFBP; INRA, Angers, France), were used to empirically find which strains were useful in defining resistance differences among aspen cultivars. For maximum diversity, the tested strains included three species (*A. tumefaciens*, *A. rhizogenes*, or *A. vitis* [formerly named biovar 1, 2, 3, respectively; 4,32,40]) harboring Ti plasmids of different opine types. Strains were isolated from various hosts (Table 1). All the strains were tumorigenic after inoculations of five herbaceous plant species commonly used for this purpose. The tumorigenicity tests with the five herbaceous plant species showed that strains had different host ranges. Oncogenic mutants were also used to show whether



*iaa* and *ipt* genes were required to define cultivar host range differences. These mutants had the chromosomal background of C58 (strain GV3101 [50]) and harbored derivatives of pTiB6S3, the Ti plasmid of the wild-type strain B6 (Table 1). They were a gift of J. P. Hernalsteens, State University, Brussels, and A. Reynaerts, Plant Genetic System, Gent, Belgium. Plate cultures were started from single colonies from freshly revived lyophilized cultures.

**Plant materials.** Aspen cultivars used in this study were closely related hybrids of *P. tremula* L. × *P. alba* L., obtained by Lemoine (21) (Table 2). The cultivars were chosen because they

showed markedly different responses to natural infections (30). Also, 'Astria', a triploid hybrid that is intensively used for forestation in Germany, was included. The reliability of the resistance assessment was tested with aspens propagated either in vitro or in growth chambers under a 16 h of light/8 h of dark photoperiod, with light provided by fluorescent tubes at a photon flux density of 50  $\mu\text{E m}^{-2} \text{s}^{-1}$ . Different temperature regimes ( $22 \pm 1^\circ\text{C}$  in vitro,  $25 \pm 1^\circ\text{C}$  or  $19 \pm 1^\circ\text{C}$  in growth chambers) were used, since this parameter can affect agrobacterial virulence (9). Shoot cultures were established on Murashige and Skoog (MS) medium (28) with half-concentrated mineral salts.

TABLE 1. Characteristics of *Agrobacterium* spp. strains used in virulence assays on aspen cultivars

Wild-type strain <sup>a</sup>		Host of origin	Species of <i>Agrobacterium</i> <sup>b</sup>	Opine type of the pTi <sup>c</sup>	Tumorigenicity <sup>d</sup>
CFBP no.	Other code				
Virulence group C58 <sup>e</sup>					
1903	C58	<i>Prunus cerasus</i>	<i>tumefaciens</i>	Nopaline	5
296	111	Unknown	<i>tumefaciens</i>	Nopaline	5
449		<i>Prunus domestica</i>	<i>rhizogenes</i>	Agropine	5
1317		<i>Rubus idaeus</i>	<i>rhizogenes</i>	Nopaline	5
1804		<i>Prunus persicae</i>	<i>rhizogenes</i>	Nopaline	5
1840		<i>Populus alba</i>	<i>rhizogenes</i>	Nopaline	5
1873		<i>Dahlia</i> sp.	<i>tumefaciens</i>	Agropine	5
1932		<i>Prunus persicae</i>	<i>tumefaciens</i>	Nopaline	5
1933		<i>Malus pumila</i>	<i>tumefaciens</i>	Nopaline	5
1961		<i>Populus bolleana</i>	<i>rhizogenes</i>	Nopaline	5
1962		<i>Prunus mahaleb</i>	<i>rhizogenes</i>	Nopaline	5
2177		<i>Populus tremula</i> × <i>P. alba</i>	<i>tumefaciens</i>	Nopaline	5
2178	82-139	<i>Prunus avium</i>	<i>rhizogenes</i>	Nopaline	5
2326		<i>Prunus avium</i>	<i>rhizogenes</i>	Nopaline	5
2410	M22	<i>Populus</i> sp.	<i>tumefaciens</i>	Nopaline	5
2411	CG4	<i>Salix purpurea</i>	<i>tumefaciens</i>	Nopaline	5
2516		<i>Populus tremula</i> × <i>P. alba</i>	<i>tumefaciens</i>	Nopaline	5
	Bo542	<i>Dahlia</i> sp.	<i>tumefaciens</i>	Agropine	5
Virulence group B6					
2413	B6	<i>Malus</i> sp.	<i>tumefaciens</i>	Octopine	5
1935		<i>Rosa</i> sp.	<i>rhizogenes</i>	Nopaline	4
Virulence group 1904					
1904	AG20	<i>Vitis vinifera</i>	<i>tumefaciens</i>	Nopaline	2
1905	AG28	<i>Vitis vinifera</i>	<i>rhizogenes</i>	Nopaline	3
2179		<i>Vitis vinifera</i>	<i>vitis</i>	Nopaline	3
Virulence group 354					
354	B10	Unknown	<i>tumefaciens</i>	Nopaline	1
450		<i>Malus pumila</i>	<i>rhizogenes</i>	Unknown	3
452		<i>Malus pumila</i>	<i>tumefaciens</i>	Unknown	3
1930		<i>Malus pumila</i>	<i>rhizogenes</i>	Unknown	2
2412	A1B6	Unknown	<i>tumefaciens</i>	Octopine	3
Virulence group 1001					
1001	ATCC 13333	<i>Malus sylvestris</i>	<i>tumefaciens</i>	Octopine	4
2407		<i>Vitis vinifera</i>	<i>tumefaciens</i>	Octopine cucumopine	2

Mutant strain<sup>f</sup>

Chromosomal background	Relevant characters	Reference
GV3101	Derivative of C58 cured of pTiC58	50
Plasmid		
pTiB6S3	Nonmutated control	
pGV2250	Insertion mutant in the <i>ipt</i> gene	35
pGV2255	Deletion of <i>ipt</i> , 6a and 6b	29
pGV2215	<i>iaaM</i> and <i>iaaH</i> genes deleted	20
pGV2235	Only <i>ipt</i> , <i>ocs</i> , and <i>Km<sup>r</sup></i> in the T-DNA	20
pGV2282	Insertion mutant in the <i>iaaM</i> gene	20
pGV2274	Insertion mutant in the <i>iaaH</i> gene	17

<sup>a</sup> Strains were mainly collected by Collection Française de Bactéries Phytopathogènes (CFBP) collaborators (Ridé, Lopez, Prunier, Nesme, and Michel), otherwise the usual codes have been indicated.

<sup>b</sup> *Agrobacterium* species correspond to biovars.

<sup>c</sup> The pTi opine types were determined according to Petit et al. (34).

<sup>d</sup> Tumorigenicity indicates the number of plant species that formed tumors when the strain was inoculated to tomato, tobacco, *Datura stramonium* and *Kalanchoë tubiflora* seedlings, and carrot root disks.

<sup>e</sup> Virulence groups were determined in vitro with five selected aspen cultivars.

<sup>f</sup> *ipt* encodes cytokinin biosynthesis; *iaaM* and *iaaH* are required together for auxin biosynthesis; 6a is required for opine excretion; 6b is required for tumor morphology; and *ocs* is required for octopine biosynthesis. *Km<sup>r</sup>* indicates resistance to kanamycin.



**Inoculation procedures.** Inoculations were performed in wounded, subapical, nonlignous internodes. For aspens generated in vitro, wounds were made with a needle and inoculated with a dense bacterial suspension optically adjusted to  $3 \times 10^8$  cells ml<sup>-1</sup>. Aspens cultivated in growth chambers or herbaceous plants grown in the greenhouse were pricked with a toothpick previously dipped in plate cultures. Carrot root disks were cultivated axenically on water agar, and inoculated with a dense bacterial suspension. Each strain-cultivar combination was tested with four plants inoculated at two to three sites, and each trial was replicated twice with the wild-type strains and three times with the mutants. Control inoculations were performed with sterile distilled water. Opines were detected in tumors according to the procedure of Petit et al. (34).

**Percentage of crown gall.** Percentages of crown galled plants in a naturally infected nursery were determined by visual inspection of aspen root systems once in 1984 and once in 1986. These two instances corresponded to cutting harvests of two successive campaigns of multiplication. The 1984/1986 multiplication campaign was performed with root material obtained from the previous 1982/1984 campaign, and used only healthy material selected after a drastic culling of the diseased material. Aspens were propagated in the nursery by the root cutting method (30).

**Sensitivity to cytokinin.** A leaf disk assay was used to estimate cultivar sensitivity to cytokinin. The fourth and fifth leaves beneath the apexes of aspens cultivated in growth chambers or in the greenhouse were surface-sterilized with 4 mM HgCl<sub>2</sub> for 20 min, rinsed twice in 17 mM CaCl<sub>2</sub> for 10 min, and then incubated for 1 h in sterile water. Disks, 8-mm in diameter, were punched between major veins and randomly plated on standard MS medium supplemented with carbenicillin at 500 mg liter<sup>-1</sup>, Benlate (benomyl; du Pont, Boston) at 100 mg liter<sup>-1</sup>, and BA (benzyladenine; Sigma Chemical Co., St. Louis) at 0.03 to 32 μM. BA solutions were obtained by twofold serial dilutions, and then they were filter-sterilized and added to MS medium after autoclaving. Sixteen to thirty disks per cultivar-BA concentration combination were used in two independent experiments. Sealed plates were incubated at 22 ± 1°C under a 16 h light/8 h dark photoperiod, with light provided by fluorescent tubes at a photon flux density of 50 μE m<sup>-2</sup> s<sup>-1</sup>. Leaf disks were scored for morphological changes twice a week for 3 months. The sensitivity to exogenous cytokinin was estimated by calculating the ED<sub>50</sub> value, defined as the BA concentration causing necrosis on 50% of disks after 40 to 45 days.

**Data analysis.** Statistical analyses were performed according to Snedecor and Cochran (44) using the Stat View 4.0 software (Abacus Concepts, Inc., Berkeley, CA).

## RESULTS

**Resistance assessment of aspen cultivars using wild-type *Agrobacterium* spp.** In initial experiments, strains of *Agrobacterium* spp. isolated from aspen crown galls (i.e., 1961, 2516, 2177, 1840, and 2410) formed tumors with every tested aspen cultivar. Therefore, a larger set of strains was screened to select those that could define differences in cultivar resistance in a reliable manner. Virulence screening of the 30 strains was performed with plants generated in vitro, because tumor formation is optimal in this condition. Since the manipulation of a large number of cultivars in vitro is difficult, five aspen cultivars were selected based on differences in susceptibility to natural infections. The 30 wild-type strains were divided into five groups of identical virulence, allowing for discrimination of four resistance groups among the five selected aspen cultivars (Table 3). The reliability of the resistance assessment of the five cultivars was tested in growth chambers and in vitro at various temperature regimes (19, 25, and 22°C, respectively) with virulence group representatives C58, B6, 1904, and 354. Reliable results were obtained with C58, B6, and

354, but not with 1904, which was very sensitive to temperature (data not shown). Therefore, the resistance of the remaining aspen cultivars was estimated in a growth chamber at 19°C with the three selected strains that gave reliable results. All aspen cultivars could then be ranked into three classes of resistance: susceptible, intermediate, and resistant (Table 4).

**Relation with response to natural infections.** The response to natural infections was estimated by the percentage of crown galled plants per cultivar determined in two instances (Table 5). Data were compared to the resistance rank of aspen defined by artificial infections, and a significant rank correlation was found (Kendall's rank correlation corrected for ties,  $P = 0.016$ ). The Kendall's coefficient of correlation ( $\tau = -0.301$ ) was rather small, however, suggesting that only a small part of the disease response was controlled by differential plant susceptibility to *A. tumefaciens*. Nevertheless, the most salient feature was that resistant aspens (did not form tumors with B6) consistently had the lowest percentage of crown galls (mean percentage = 4.5%). As a result, resistance to B6 could be considered as an indicator of the tolerance of aspen cultivars to crown gall under natural conditions.

**Assays with T-DNA mutants of pTiB6S3.** The following experiment was aimed at defining whether T-DNA oncogenes of B6 were involved in differences in susceptibility among aspen cultivars. For this purpose, aspen cultivars (selected for their differences in susceptibility to artificial infection) were inoculated with a strain harboring various derivatives of the Ti plasmid of B6 that were either *iaa*- or *ipt*-defective. Results showed that the aspen cultivars had different resistance patterns when inoculated with

TABLE 2. Pedigrees of the aspen (*Populus* spp.) cultivars used to find differences in susceptibility to crown gall

Cultivar <sup>a</sup>	Parent	
	<i>P. tremula</i> (female)	<i>P. alba</i> (male)
706-1	5820	5872
706-2	5820	5872
706-3	5820	5872
706-8	5820	5872
709-1	5813	5872
709-27	5813	5872
709-3	5813	5872
710-23	5814	5872
710-24	5814	5872
710-27	5814	5872
712-1	5815	5872
712-7	5815	5872
712-8	5815	5872
717-1-1 <sup>b</sup>	5903	6072
717-1-2	5903	6072
717-1-B4	5903	6072
'Austria'	Unknown	<i>P. tremuloides</i>

<sup>a</sup> Cultivars were F<sub>1</sub> hybrids obtained from wild-type progenitors.

<sup>b</sup> The 717-1-1 hybrid is registered in the European list of poplar cultivars authorized for plantation under the name 'Rajane.'

TABLE 3. Resistance patterns of selected in vitro generated aspen plants to wild-type strains of *Agrobacterium* spp.

Cultivar	Virulence group <sup>a</sup>				
	C58	B6	1904	354	1001
717-1-B4	+ <sup>b</sup>	+	+	+	-
712-7	+	+	+	+	-
710-23	+	+	+/-	-	-
709-3	+	+	-	-	-
710-24	+	-	-	-	-

<sup>a</sup> Tests were performed at 22°C with 30 wild-type strains in two independent experiments. Strains with identical virulence were grouped in virulence groups as described in Table 1.

<sup>b</sup> + indicates that 29 to 100% of inoculated sites formed tumors after 1 month; +/- indicates 8% formed tumors; and - indicates no tumors formed.



oncogene mutants (Table 6). Therefore, since mutant bacteria had the same bacterial background (i.e., strain GV3101), the bacterial ability to define resistance differences among aspen cultivars appeared to be mainly determined by the Ti plasmid, and especially T-DNA oncogenes.

Among the T-DNA oncogenes, the various *iaa*-defective mutants formed tumors with cultivar 710-23, but not with cultivars 709-3 or 710-24. This showed that *iaa* genes were not absolutely

TABLE 4. Resistance assessment of aspen cultivars with selected strains of *Agrobacterium tumefaciens*

Cultivar	<i>A. tumefaciens</i> strain		
	C58	B6	354
Susceptible			
717-1-B4	+ <sup>a</sup>	+	+
712-7	+	+	+
706-2	+	+	+
706-3	+	+	+
'Astria'	+	+	+
Intermediate			
709-3	+	+	-
710-23	+	+	-
709-27	+	+	-
709-1	+	+	-
712-8	+	+	-
706-8	+	+	-
706-1	+	+	-
Resistant			
710-24	+	-	-
712-1	+	-	-
710-27	+	-	-
717-1-1	+	-	-
717-1-2	+	-	-

<sup>a</sup> Tests were performed in a growth chamber at 19°C in two independent experiments. + indicates the presence of tumors containing expected opines, and - indicates the absence of both tumors and opine at inoculation sites.

TABLE 5. Percentage of crown gall in aspen cultivars in a naturally infected nursery

Cultivar	Multiplication campaign	
	1982/1984	1984/1986
Susceptible <sup>a</sup>		
717-1-B4	42 / 160 <sup>b</sup>	86 / 497
712-7	0 / 118	10 / 428
706-2	97 / 170	150 / 381
706-3	13 / 199	70 / 530
Mean percentage <sup>c</sup>	16%	
Intermediate		
709-3	102 / 108	9 / 18
710-23	4 / 75	13 / 198
709-27	19 / 58	7 / 181
709-1	27 / 173	16 / 142
712-8	17 / 191	3 / 363
706-8	45 / 171	44 / 241
706-1	132 / 189	31 / 239
Mean percentage	22%	
Resistant		
710-24	4 / 149	1 / 142
712-1	2 / 177	1 / 422
710-27	2 / 103	16 / 336
717-1-1	21 / 136	35 / 401
717-1-2	31 / 192	31 / 500
Mean percentage	4.5%	

<sup>a</sup> The classes susceptible, intermediate, and resistant were as defined in Table 4.

<sup>b</sup> Data indicates the number of crown galled plants out of the total number of plants assayed.

<sup>c</sup> Mean percentages calculated after arcsin  $\sqrt{p}$  transformation.

required to define resistance differences. On the other hand, mutants defective for cytokinin biosynthesis formed tumors and roots with all cultivars, including 710-24, which was resistant to the unmutated parental strain. Thus, a defective *ipt* gene did not abolish the tumorigenicity of strains harboring a pTiB6S3, but abolished the ability to define resistance differences between aspens. This suggested a role of the sensitivity to the *ipt* gene product, a cytokinin, in the individual plant ability to form tumors.

**Cytokinin sensitivity of aspen cultivars.** An experiment was attempted to define intrinsic differences in sensitivity to exogenous cytokinin of aspen cultivars. For this purpose, aspen leaf disks were incubated on media containing various amounts of BA, a cytokinin. On BA-free medium, the leaf disks survived 60 to 70 days; they then became chlorotic and degenerated slowly without browning of tissues. On BA-supplemented media, disks turned brown after 30 days and died a week later. Striking differences in sensitivity were observed among aspen cultivars (Fig. 1), allowing for the determination of ED<sub>50</sub> values, defined as the BA concentration that kills 50% of disks after 40 to 45 days (Table 7). Observations showed that aspens with low ED<sub>50</sub> values often developed 1- to 4-mm calluses that, in most cases, later gave rise to roots on BA-free medium or on medium containing <1 μM BA (Mann-Whitney's test for callus production effect on ED<sub>50</sub> corrected for ties, *P* = 0.036).

**Relationship between sensitivity to cytokinin and susceptibility to crown gall.** The susceptibility to *A. tumefaciens* strains could largely be explained by the sensitivity to cytokinin as estimated by ED<sub>50</sub> values (Table 8). The only exception was cultivar 706-3 (Table 7). Callus production was also significantly related to the susceptibility to *A. tumefaciens* strains (Table 8), with the exception of cultivar 717-1-2 (Table 7).

The percentage of plants crown galled in a naturally contaminated nursery could also be explained by the ED<sub>50</sub> values and by callus production (Table 8). In this case, the value of the Kendall's coefficient ( $\tau$  = -0.31) was rather small, suggesting that only a small part of the susceptibility to galling was controlled by the plant sensitivity to cytokinin.

## DISCUSSION

In spite of its economic importance for many crops and extensive studies on genetic control, attempts to find intrinsic plant factors involved in the resistance to crown gall are rather rare. Cultivars resistant to *A. tumefaciens* have been described in several host plants including *Nicotiana* spp. (49), *Arabidopsis thaliana* (22), *Cucurbita maxima* (43), *Pisum sativum* (39), *Populus deltoides* (38), *Glycine max* (33), *Medicago sativa* (24), *Prunus* spp. (36), *Rubus idaeus* (54), *Chrysanthemum morifolium* (6,26), *Vitis* spp. (47), and *Rosa* spp. (3). In this list, only the latter six crops significantly suffer from crown gall outbreaks, and, even in these cases, there is no information about resistance of the se-

TABLE 6. Virulence of mutated *Agrobacterium tumefaciens* strains on aspen cultivars

Cultivar <sup>a</sup>	Plasmid content of strain GV3101 <sup>b</sup>		
	<i>ipt</i> -deficient mutants (pGV2250, pGV2255)	Unmutated control (pTiB6S3)	<i>iaa</i> -deficient mutants (pGV2215, pGV2235, pGV2282, pGV2274)
710-23	+ r	+	+ b
709-3	+ r	+	-
710-24	+ r	-	-

<sup>a</sup> Assays were performed with plants generated in vitro at 22°C in three independent experiments.

<sup>b</sup> + indicates tumors; - indicates no tumors formed; b indicates the presence of buds associated with tumors formed by a strain harboring pGV2215; r indicates the presence of roots associated with tumors with the strains harboring pGV2250 and pGV2255.



lected material in natural conditions. Moreover, the study of plant factors involved in resistance often utilizes model host plants, which are generally not used agronomically. To our knowledge, no work has yet established a link between crown gall in experimental plots naturally contaminated by the pathogen with a resistance assessment done after artificial infections and the finding of an intrinsic plant factor correlated to crown gall susceptibility. Nevertheless, all these works suggest that resistant or tolerant cultivars can also be defined in other plant species for which crown gall is significantly important. This, in turn, supports the feasibility of using genetic control under conditions in which correlations can be established between artificial infection data and behavior under natural conditions, as long as the selection for resistance does not counterselect undesirable plant traits.

*Agrobacterium* spp. strains for use in reliable resistance tests were selected with the understanding that growth conditions and temperature affect agrobacterial virulence (9). Some bacterial strains were found to be highly sensitive to temperature for unknown reasons, and these bacteria were not used to define aspen resistance. On the other hand, there was no difference in susceptibility between plants generated in vitro and plants cultivated in growth chambers under any of the temperature regimes. Thus, the resistance of numerous aspen cultivars could be routinely defined in growth chambers with the three selected strains: C58, B6, and 354.

Among the selected strains, strain B6, which has been used to define resistance differences in several other species of host plants (6,26,39,43), has a particular agronomic interest, since aspen cultivars resistant to this strain consistently had the lowest rate of crown gall in naturally contaminated plots. Nevertheless, the field resistance obtained by a selection of cultivars resistant to B6 was not enough to get an acceptable control of the disease without sanitation practices. A drastic culling of the diseased material (30), or preferably the use of *Agrobacterium*-free plants generated in vitro (C. Ponsonnet and X. Nesme, unpublished data), must also be implemented to reach a crown gall percentage lower than 5 to 10%, a maximum acceptable level for a disease affecting plant quality (2). A higher level of resistance in aspen cultivars is, therefore, desirable. Resistance to C58 could serve as an indicator, since poplars of other sections which are generally not affected by crown gall are generally resistant to this strain (38).

Bacterial genes involved in defining resistance differences were studied using strain B6. It has been hypothesized that host-*Agrobacterium* incompatibility could be because of the inability of the bacteria to bind host cells (25), noninduction of virulence genes

(41), or induction of a hypersensitive response caused by toxic overproduction of phytohormones at the inoculation sites by both *vir* and T-DNA genes (53). These various hypotheses were tested by inoculating mutant agrobacteria to aspens. As in several other systems (19,23,48), incompatibility to B6 appeared to be mainly determined by its Ti plasmid. Thus, it was unlikely that *Agrobacterium*-host cell binding was involved in aspen resistance, since this step is controlled by chromosomal genes (8,13). The incompatibility also does not result from a lack of induction or a toxic effect of *vir* genes, since one cultivar resistant to a strain harboring pTiB6S3 formed tumors with other strains carrying T-DNA mutants of the same plasmid having identical *vir* genes. On the other hand, T-DNA oncogenes, especially *ipt*, appeared to be strongly involved, since mutants retained their ability to define resistance differences as long as they had a functional *ipt* gene, and this property was abolished in the absence of this gene. Involvement of *ipt* in host range definitions has been described in the *A. vitis*/grapevine system. In this system, a functional gene from a virulent strain permitted expansion of the host range of the limited host range (LHR) strains which had nonfunctional *ipt* (6,16). It is not known, however, whether virulence differences among 354, B6, and C58 were because of differential expressions of their respective *ipt* genes.

Our results strongly implied that hormone sensitivity could control susceptibility to crown gall in aspens. Thus, a leaf disk

TABLE 7. Sensitivity to exogenous cytokinin of aspen cultivars

Cultivar	Sensitivity to cytokinin <sup>a</sup>	
	ED <sub>50</sub> <sup>b</sup>	Callus production <sup>c</sup>
Susceptible <sup>d</sup>		
717-1-B4	2	+
712-7	2	+
706-3	16	+
'Astria'	2	+
Intermediate		
709-3	4	+
710-23	4	+
712-8	8	-
706-8	8	+
Resistant		
710-24	32	-
712-1	32	-
710-27	32	-
717-1-2	32	+

<sup>a</sup> Determined by a leaf disk assay in two independent experiments.

<sup>b</sup> The benzyladenine (BA) concentration killed 50% of leaf disks after 45 to 50 days.

<sup>c</sup> Callus production after 45 to 50 days; + indicates the presence of calli on leaf disks plated below 1  $\mu$ M BA, generally developing into roots later; and - indicates no callus.

<sup>d</sup> Classes of resistance to *A. tumefaciens* strains are defined in Table 4.

TABLE 8. Correlation between sensitivity to cytokinin and susceptibility to crown gall of aspen cultivars<sup>a</sup>

Sensitivity to cytokinin	Susceptibility to selected strains of <i>A. tumefaciens</i> C58, B6, 354	Percentage of crown galled plants per clone in a naturally contaminated nursery
	ED <sub>50</sub>	$n = 12$ $\tau = 0.78$ ( $\tau^2 = 60\%$ ) $P = 0.0004$
Callus production	$n = 14$ $P = 0.020$	$n = 26$ $P = 0.002$

<sup>a</sup> Effects of the variables ED<sub>50</sub> value and callus production on aspen susceptibility to crown gall were tested by Kendall's rank of correlation and the Mann-Whitney *U* test, respectively, after eventual corrections for ties. Detailed data are available in Tables 4, 5, and 7.




Fig. 1. Effect of the amount of exogenous benzyladenine (BA) on the survival of leaf disks in different aspen cultivars. Sensitivity assays were performed in vitro at 22°C with leaf disks plated on Murashige and Skoog medium containing various amounts of BA in two independent experiments. Data were scored after 40 to 45 days.



assay was developed to measure cultivar sensitivity to exogenous cytokinin. Individual responses of closely related cultivars varied from 2 to 32  $\mu$ M BA. We do not know, however, whether toxicity (which defines ED<sub>50</sub> value) was a real property of cytokinins as it is for auxins (7), or if it was in a phenotype expressed only by surviving explants. ED<sub>50</sub> values showed that resistant aspens were not very sensitive to cytokinin. Thus, our hypothesis was confirmed, even if exceptions to the rule indicated that other unknown plant factors must also control the phenomenon. Plant genotypes resistant to auxin are resistant to *A. tumefaciens* strains (15,22,49). However, to our knowledge, the involvement of the sensitivity to cytokinin in the control of susceptibility to crown gall has been hypothesized, but not clearly established. As suspected in other systems, resistant aspens seemed to require more cytokinin than susceptible cultivars to reach the cytokinin/auxin ratio required for tumor development (18).

This study showed that individual resistance to *A. tumefaciens* of aspen cultivars was largely related to hormonal sensitivity. This relationship probably occurs in many other host plants susceptible to crown gall. Studies of this relationship could facilitate the study of plant contribution to crown gall development, including, for instance, the host-mediated constraints to plant transformation with *A. tumefaciens*.

#### LITERATURE CITED

- Akiyoshi, D. E., Klee, H., Amasino, R. M., Nester, E. W., and Gordon, M. P. 1984. T-DNA of *Agrobacterium tumefaciens* encodes an enzyme of cytokinin biosynthesis. Proc. Natl. Acad. Sci. USA 81:5994-5998.
- Anonymous. 1993. Liste d'organismes spécifiques nuisibles quant à la qualité selon les différents genres et espèces. J. Off. Commun. Eur. L250:4-8.
- Boelema, B. H. 1969. Resistance of rose rootstocks to crown gall (*Agrobacterium tumefaciens*). Neth. J. Plant Pathol. 75:147-150.
- Bouzar, H. 1994. Request for a judicial opinion concerning the type species of *Agrobacterium*. Int. J. Syst. Bacteriol. 44:373-374.
- Bucholz, W. G., Thomashow, M. F. 1984. Host range encoded by the *Agrobacterium tumefaciens* tumor-inducing plasmid pTiAg63 can be expanded by modification of its T-DNA oncogene complement. J. Bacteriol. 160:327-332.
- Bush, A., and Pueppke, S. 1991. Cultivar-strain specificity between *Chrysanthemum morifolium* and *Agrobacterium tumefaciens*. Physiol. Mol. Plant Pathol. 39:309-323.
- Caboche, M., Miller, J. F., Chanut, F., Aranda, G., and Cirakoglu, S. 1987. Comparison of the growth-promoting activities and toxicities of various auxin analogs on cells derived from wild type and a nonrooting mutant of tobacco. Plant Physiol. 83:795-800.
- Cangelosi, G. A., Martinetti, G., Leigh, J. A., Lee, C. C., Theines, C., and Nester, E. W. 1989. Role of *Agrobacterium tumefaciens* ChvA protein in export of  $\beta$ -1,2-glucan. J. Bacteriol. 171:1609-1615.
- Charest, P. J., and Dion, P. 1985. The influence of temperature on tumorigenesis induced by various strains of *Agrobacterium tumefaciens*. Can. J. Bot. 63:1160-1167.
- Chilton, M.-D., Drummond, H. J., Merlo, D. J., Sciaky, D., Montoya, A. L., Gordon, M. P., and Nester, E. 1977. Stable incorporation of plasmid DNA into higher plant cells: The molecular basis of crown gall tumorigenesis. Cell 11:263-271.
- De Cleene, M. 1988. The susceptibility of plants to *Agrobacterium*: A discussion of the role of phenolic compounds. FEMS (Fed. Eur. Microbiol. Soc.) Microbiol. Rev. 54:1-8.
- De Cleene, M., and De Ley, J. 1976. The host range of crown gall. Bot. Rev. 42:389-466.
- Douglas, C. J., Staneloni, R. J., Rubin, R. A., and Nester, E. W. 1985. Identification and genetic analysis of an *Agrobacterium tumefaciens* chromosomal virulence region. J. Bacteriol. 161:850-860.
- El-Fiki, F., and Giles, K. L. 1981. *Agrobacterium tumefaciens* in agriculture and research. Int. Rev. Cytol. Suppl. 13:47-58.
- Hansen, C. E., and Meins, F. 1986. Evidence for a cellular gene with potential oncogenic activity in plants. Proc. Natl. Acad. Sci. USA 83:2492-2495.
- Hoekema, A., De Pater, B. S., Fellingner, A. J., Hooykaas, P. J. J., and Schilperoort, R. A. 1984. The limited host range of *Agrobacterium tumefaciens* strain extended by a cytokinin gene from a wide host range T-region. EMBO (Eur. Mol. Biol. Organ.) J. 3:3043-3047.
- Inzé, D., Follin, A., Van Lijsebettens, M., Simoens, C., Genetello, C., Van Montagu, M., and Schell, J. 1984. Genetic analysis of the individual T-DNA genes of *Agrobacterium tumefaciens*; further evidence that two genes involved in indole-3-acid synthesis. Mol. Gen. Genet. 194:265-274.
- Ishikawa, K., Kamada, H., Yamaguchi, I., Takahashi, N., and Harada, H. 1988. Morphology and hormone levels of tobacco and carrot tissues transformed by *Agrobacterium tumefaciens*. I. Auxin and cytokinin contents of cultured tissues transformed with wild-type and mutant Ti plasmids. Plant Cell Physiol. 29:461-466.
- Knauf, V. C., Panagopoulos, C. G., and Nester, E. W. 1982. Genetic factors controlling the host range of *Agrobacterium tumefaciens*. Phytopathology 72:1545-1549.
- Leemans, J., Hernalsteens, J. P., Deblaere, R., De Greve, H., Thia-Toong, L., Van Montagu, M., and Schell, J. 1983. Genetics analysis of T-DNA and regeneration of transformed plants. Pages 322-330 in: Molecular Genetics of the Bacteria-Plant Interaction. A. Pühler, ed. Springer-Verlag KG, Berlin.
- Lemoine, M. 1973. Amélioration des peupliers de la section *Leuce* sur sols hydromorphes. Ph.D. thesis. Université de Nancy I, Nancy, France.
- Lincoln, C., Turner, J., and Estelle, M. 1992. Hormone-resistant mutants of *Arabidopsis* have an attenuated response to *Agrobacterium* strains. Plant Physiol. 98:979-983.
- Loper, J. E., and Kado, C. I. 1979. Host range conferred by the virulence-specifying plasmid of *Agrobacterium tumefaciens*. J. Bacteriol. 139:591-596.
- Mariotti, D., Davey, M. R., Draper, J., Freeman, J. P., and Cocking, E. C. 1984. Crown gall tumorigenesis in the forage legume *Medicago sativa* L. Plant Cell Physiol. 25:473-482.
- Matthysse, A. G., and Gurlitz, H. G. 1982. Plant cell range for attachment of *Agrobacterium tumefaciens* to tissue culture cells. Physiol. Plant Pathol. 21:381-387.
- Miller, H. N., Miller, J. W., and Crane, G. L. 1975. Relative susceptibility of *Chrysanthemum* cultivars to *Agrobacterium*. Plant Dis. Rep. 59:576-581.
- Moore, L. W., and Cooksey, D. A. 1981. Biology of *Agrobacterium tumefaciens*: Plant interactions. Int. Rev. Cytol. Suppl. 13:15-46.
- Murashige, T., and Skoog, F. 1962. A revised medium for rapid growth and bioassays with tobacco tissue cultures. Physiol. Plant. 15:473-497.
- Nacmias, B., Ugolini, S., Ricci, M. D., Pellegrini, M. G., Bogani, P., Bettini, P., Inzé, D., and Buiatti, M. 1987. Tumor formation and morphogenesis on different *Nicotiana* sp. and hybrids induced by *Agrobacterium tumefaciens* T-DNA mutants. Dev. Genet. 8:61-71.
- Nesme, X., Beneddra, T., and Collin, E. 1990. Importance du crown gall chez les hybrides de *Populus tremula* L.  $\times$  *P. alba* L. en pépinière forestière. Agronomie 10:581-583.
- Nesme, X., Michel, M.-F., and Digat, B. 1987. Population heterogeneity of *Agrobacterium tumefaciens* in galls of *Populus* L. from a single nursery. Appl. Environ. Microbiol. 53:655-659.
- Ophel, K., and Kerr, A. 1990. *Agrobacterium vitis* sp. nov. for strains of *Agrobacterium* biovar 3 from grapevines. Int. J. Syst. Bacteriol. 40:236-241.
- Owens, L., and Cress, D. E. 1985. Genotypic variability of soybean response to *Agrobacterium* strains harboring the Ti or Ri plasmids. Plant Physiol. 77:87-94.
- Petit, A., Berkaloff, A., and Tempé, J. 1986. Multiple transformation of plant cell by *Agrobacterium* may be responsible for the complex organization of T-DNA in crown gall and hairy root. Mol. Gen. Genet. 202:388-393.
- Petit, A., Tempé, J., Kerr, A., Holsters, M., Van Montagu, M., and Schell, J. 1978. Substrate induction of conjugative activity of *Agrobacterium tumefaciens* Ti plasmids. Nature (Lond.) 271:570-572.
- Pierronnet, A., and Escalettes, V. 1990. Study on the behaviour of *Prunus* cultivars towards *Agrobacterium radiobacter* var. *tumefaciens*. Acta Hort. (Wageningen) 283:327-332.
- Ream, W. 1989. *Agrobacterium tumefaciens* and interkingdom genetic exchange. Annu. Rev. Phytopathol. 27:583-618.
- Riemenschneider, D. E. 1990. Susceptibility of infra- and inter-specific hybrid poplars to *Agrobacterium tumefaciens* strain C58. Phytopathology 80:1099-1102.
- Robbs, S. T., Hawes, M. C., Lin, H. J., Pueppke, S. G., and Smith, L. Y. 1991. Inheritance of resistance to crown gall in *Pisum sativum*. Plant Physiol. 95:52-57.
- Sawada, H., Ieki, H., Oyaizu, H., and Matsumoto, S. 1993. Proposal for rejection of *Agrobacterium tumefaciens* and revised descriptions for the genus *Agrobacterium* and for *Agrobacterium radiobacter* and *Agrobacterium rhizogenes*. Int. J. Syst. Bacteriol. 43:694-702.
- Schafer, W., Görz, A., and Kahl, G. 1987. T-DNA integration and expression in a monocot crop plant after induction of *Agrobacterium*. Nature (Lond.) 327:529-532.



- E 9 9 A 0 0 0
42. Schröder, G., Klipp, W., Hillebrand, A., Ehring, R., Koncz, C., and Schröder, J. 1983. The conserved part of the T-region in Ti-plasmids expresses four proteins in bacteria. *EMBO (Eur. Mol. Biol. Organ.) J.* 2:403-409.
  43. Smarelli, J., Watters, M. T., and Diba, L. H. 1986. Response of various cucurbits to infection by plasmid-harboring strains of *Agrobacterium*. *Plant Physiol.* 82:622-624.
  44. Snedecor, G. W., and Cochran, W. G. 1957. *Statistical Methods*, 6e ed. Iowa State University Press, Ames.
  45. Stachel, S. E., Nester, E., and Zambryski, P. 1986. A plant cell factor induces *Agrobacterium tumefaciens vir* gene expression. *Proc. Natl. Acad. Sci. USA* 83:379-383.
  46. Stomp, A.-M., Loopstra, C., Chilton, W., Sederoff, R., and Moore, L. 1990. Extended host range of *Agrobacterium tumefaciens* in the genus *Pinus*. *Plant Physiol.* 92:1226-1232.
  47. Süle, S., Mozsar, J., and Burr, T. J. 1994. Crown gall resistance of *Vitis* spp. and grapevine rootstocks. *Phytopathology* 84:607-611.
  48. Thomashow, M., Panagopoulos, C., Gordon, M., and Nester, E. 1980. Host range of *Agrobacterium tumefaciens* is determined by the Ti plasmid. *Nature (Lond.)* 283:794-796.
  49. Tourneur, J., Jouanin, L., Muller, J.-F., Caboche, M., and Simpson, R. 1985. A genetic approach to study the mechanism of action of auxin in tobacco. Susceptibility of an auxin resistant mutant to *Agrobacterium* transformation. Pages 791-797 in: *Plant Genetics*. M. Freeling, ed. Alan R. Liss, Inc., New York.
  50. Van Larebecke, N., Englers, G., Holters, M., Van Den Elsacker, S., and Zaenen, I. 1974. Large plasmid in *Agrobacterium tumefaciens* is essential for crown gall-inducing ability. *Nature (Lond.)* 252:169-170.
  51. Winans, S. C. 1992. Two-way chemical signaling in *Agrobacterium*-plant interactions. *Microbiol. Rev.* 56:12-31.
  52. Yanofsky, M., Montoya, A., Knauf, V., Lowe, B., Gordon, M. P., and Nester, E. W. 1985. Molecular and genetic analysis of factors controlling host range in *Agrobacterium tumefaciens*. *Mol. Gen. Genet.* 201:237-246.
  53. Yanofsky, M. F., and Nester, E. W. 1986. Molecular characterization of a host-range-determining locus from *Agrobacterium tumefaciens*. *J. Bacteriol.* 168:244-250.
  54. Zurowski, C. L., Copeman, R. J., and Daubeny, H. A. 1985. Relative susceptibility of red raspberry clones to crown gall. (Abstr.) *Phytopathology* 75:1289.