


Volume 134 No. 1 March 2005 ISSN 0269-7491



# ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

**W.J. Manning**


ASSOCIATE EDITORS

**K.E. Havens**

**S.V. Krupa**

**K.C. Jones**

**J. W. Erisman**





# ENVIRONMENTAL POLLUTION

## CONTENTS—Continued from outside back cover

- 97 **Contamination of rivers in Tianjin, China by polycyclic aromatic hydrocarbons**  
Z. Shi, S. Tao, B. Pan, W. Fan, X.C. He, Q. Zuo, S.P. Wu, B.G. Li, J. Cao, W.X. Liu, F.L. Xu, X.J. Wang, W.R. Shen, P.K. Wong  
*Coal combustion is suggested as a recent local source of PAHs in this area.*
- 113 **Bioaccumulation of PCBs in the cuttlefish *Sepia officinalis* from seawater, sediment and food pathways**  
B. Danis, P. Bustamante, O. Cotret, J.L. Teyssié, S.W. Fowler, M. Warnau  
*Bioaccumulation of PCBs by cuttlefish is studied, via seawater, sediments and their food.*
- 123 **The combined effect of lead exposure and high or low dietary calcium on health and immunocompetence in the zebra finch (*Taeniopygia guttata*)**  
T. Snoeijs, T. Dauwe, R. Pinxten, V.M. Darras, L. Arckens, M. Eens  
*Male and female finches may respond to lead differently.*
- 133 **In situ exposures using caged organisms: a multi-compartment approach to detect aquatic toxicity and bioaccumulation**  
G.A. Burton, M.S. Greenberg, C.D. Rowland, C.A. Irvine, D.R. Lavoie, J.A. Brooker, L. Moore, D.F.N. Raymer, R.A. McWilliam  
*In situ exposures provide unique information that is complementary to traditional lab-based toxicity results.*
- 145 **Using plant biomonitors and flux modelling to develop O<sub>3</sub> dose-response relationships in Catalonia**  
I. Filella, J. Peñuelas, A. Ribas  
*Modelling of biomonitors ozone absorbed dose improves damage estimation in comparison with exposure indices such as AOT40.*
- 153 **The maximum reservoir capacity of soils for persistent organic pollutants: implications for global cycling**  
M.D. Valle, E. Jurado, J. Dachs, A.J. Sweetman, K.C. Jones  
*Major global terrestrial sinks/stores for POPs are identified and the significance of gradients between them discussed.*
- 165 **Water striders (family Gerridae): mercury sentinels in small freshwater ecosystems**  
T.D. Jardine, T.A. Al, K.T.B. MacQuarrie, C.D. Ritchie, P.A. Arp, A. Maprani, R.A. Cunjak  
*Water striders accurately reflect the entry of mercury in food chains of small freshwater systems.*
- 173 **Mercury baseline levels in Flemish soils (Belgium)**  
F.M.G. Tack, T. Vanhaesebroeck, M.G. Verloo, K. Van Rompaey, E. Van Ranst  
*Increased mercury levels may reflect human activity.*



**CONTENTS**

**Volume 134 Number 1 2005**

**New initiative**

- 1 Quantifying ozone uptake and its effects on the stand level of common beech (*Fagus Sylvatica* L.) in Southern Germany**  
C. Dittmar, K. Pfaffelmoser, T. Rötzer, W. Elling

*Water balance calculations can be used to estimate long-term O<sub>3</sub> uptake at the stand level and in combination with tree-ring data to derive new critical threshold values.*

**Research papers**

- 5 Validated sampling strategy for assessing contaminants in soil stockpiles**

F. Lamé, T. Honders, G. Derksen, M. Gadella

*A sampling strategy that ensures analytical results representative of the mean concentration in soil stockpiles is presented and validated.*

- 13 Mercury transfer from fish carcasses to scavengers in boreal lakes: the use of stable isotopes of mercury**

J. Sarica, M. Amyot, L. Hare, P. Blanchfield, R.A. (Drew) Bodaly, H. Hintelmann, M. Lucotte

*Scavengers such as leeches accumulate mercury from fish carcasses which then may return into the food web.*

- 23 Study of calcium-dependent lead-tolerance on plants differing in their level of Ca-deficiency tolerance**

D.M. Antosiewicz

*Calcium regulated lead deposition in cell walls of plants.*

- 35 Sorption studies of chloroanilines on kaolinite and montmorillonite**

S. Angioi, S. Polati, M. Roz, C. Rinaudo, V. Gianotti, M.C. Gennaro

*Chloroanilines are sorbed by clay standard kaolinite and Na-montmorillonite at different extents and likely by different mechanisms.*

- 45 Bioaccumulation of <sup>51</sup>Cr, <sup>63</sup>Ni and <sup>14</sup>C in Baltic Sea benthos**

L. Kumblad, C. Bradshaw, M. Gilek

*Bioaccumulation of radioisotopes in Baltic Sea benthos has important implications for contaminant transfer and exposure.*

- 57 Do tubificid worms influence the fate of organic matter and pollutants in stormwater sediments?**

F. Mermillod-Blondin, G. Nogaro, T. Datry, F. Malard, J. Gibert

*Tubificid worms significantly increased the organic matter mineralization and the release of nutrients from stormwater sediments.*

- 71 Release behavior of triazine residues in stabilised contaminated soils**

G.G. Ying, R.S. Kookana, M. Mallavarpu

*Stabilisation of contaminated soil with a mix of activated carbon and cement may fail to immobilize some contaminants like triazines.*

- 79 Mercury contamination in human hair and fish from Cambodia: levels, specific accumulation and risk assessment**

T. Agusa, T. Kunito, H. Iwata, I. Monirith, T.S. Tana, A. Subramanian, S. Tanabe

*A source other than fish may be responsible for high Hg in some Cambodians.*

- 87 Impact of a flood disaster on sediment toxicity in a major river system – the Elbe flood 2002 as a case study**

M. Oetken, B. Stachel, M. Pfenninger, J. Oehlmann

*The extraordinary Elbe flood in August 2002 did not result in an overall increase of environmental contamination.*