

Volume 135 No. 1 May 2005 ISSN 0269-7491

ENVIRONMENTAL POLLUTION


EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens **S.V. Krupa**

K.C. Jones **J. W. Erisman**



ENVIRONMENTAL POLLUTION

CONTENTS—Continued from outside back cover

- 101 **Isoprene emission from tropical tree species**
P.K. Padhy, C.K. Varshney
Isoprene flux (diurnal and seasonal) from some tropical tree species was estimated and a regional comparison was made.
- 111 **Epiphytic lichen diversity on dead and dying conifers under different levels of atmospheric pollution**
M. Hauck
Epiphytic lichen diversity differs increasingly between healthy and dead or dying conifers with increasing atmospheric pollutant load.
- 121 **Survival of brown trout during spring flood in DOC-rich streams in northern Sweden: the effect of present acid deposition and modelled pre-industrial water quality**
H. Laudon, A.B.S. Poléo, L.A. Vøllestad, K. Bishop
High levels of complexed aluminum, at pH levels below 5.0, predisposes brown trout to sulfur-caused damage in the spring.
- 131 **Seasonal exposures to triazine and other pesticides in surface waters in the western Highveld corn-production region in South Africa**
L.H. Du Preez, P.J. Jansen van Rensburg, A.M. Jooste, J.A. Carr, J.P. Giesy, T.S. Gross, R.J. Kendall, E.E. Smith, G. Van Der Kraak, K.R. Solomon
Concentrations of atrazine in an agricultural region vary greatly and are influenced by rainfall amounts.
- 143 **Atmospheric versus biological sources of polycyclic aromatic hydrocarbons (PAHs) in a tropical rain forest environment**
M. Krauss, W. Wilcke, C. Martius, A.G. Bandeira, M.V.B. Garcia, W. Amelung
Evidence of non-pyrolytic, biogenic production of PAHs is provided.
- 155 **Evaluation of quick tests for phosphorus determination in dairy manures**
A. Lugo-Ospina, T.H. Dao, J.A. Van Kessel, J.B. Reeves
The dissolved phosphorous quick test can provide real-time information on soluble manure nutrient control.
- 163 **Trace elements in loggerhead turtles (*Caretta caretta*) from the eastern Mediterranean Sea: overview and evaluation**
M.M. Storelli, A. Storelli, R. D'Addabbo, C. Marano, R. Bruno, G.O. Marcotrigiano
Cadmium concentrations in samples from turtles may be high enough to affect their health.
- 171 **Assessment of municipal waste compost as a daily cover material for odour control at landfill sites**
C. Hurst, P. Longhurst, S. Pollard, R. Smith, B. Jefferson, J. Gronow
Practical measures to improve landfill odour control are investigated.
- 179 **Mercury assessment and evaluation of its impact on fish in the Cecina river basin (Tuscany, Italy)**
R. Scerbo, T. Ristori, B. Stefanini, S. De Ranieri, C. Barghigiani
Mercury entering water from a chlor-alkali plant near Tuscany has led to contamination of river food webs.

CONTENTS

Volume 135 Number 1 2005

Research papers

- 1 Reduction of silver solubility by humic acid and thiol ligands during acanthite (β -Ag₂S) dissolution**
A.R. Jacobson, C.E. Martínez, M. Spagnuolo, M.B. McBride, P. Baveye
The presence of humic acids and thiol ligands reduced Ag dissolution from acanthite particles with mixed-oxidation surface layers.
- 11 Bioaccumulation of PCBs in the sea urchin *Paracentrotus lividus*: seawater and food exposures to a ¹⁴C-radiolabelled congener (PCB#153)**
B. Danis, O. Cotret, J.L. Teyssié, P. Bustamante, S.W. Fowler, M. Warnau
*The sea urchin *Paracentrotus lividus* is a valuable indicator for PCB contamination.*
- 17 Rapid regional recovery from sulfate and nitrate pollution in streams of the western Czech Republic – comparison to other recovering areas**
V. Majer, P. Krám, J.B. Shanley
Acidified streams experienced unprecedented recovery from sulfate and nitrate pollution but pH and ANC decreased in alkaline streams.
- 29 Effects of increased deposition of atmospheric nitrogen on an upland moor: leaching of N species and soil solution chemistry**
M.G. Pilkington, S.J.M. Caporn, J.A. Carroll, N. Cresswell, J.A. Lee, T.W. Ashenden, S.A. Brittain, B. Reynolds, B.A. Emmett
Nitrate leaching from an upland moor podsol was significantly increased only in response to rates of N deposition in excess of 96 kg N ha⁻¹ yr⁻¹.
- 41 Risk-based approach to appraise valve closure in the clam *Corbicula fluminea* in response to waterborne metals**
C.-M. Liao, L.-J. Jou, B.-C. Chen
A model was developed to link valve closure in clams to concentrations of metals in water.
- 53 Distribution of rare earth elements in an alluvial aquifer affected by acid mine drainage: the Guadiamar aquifer (SW Spain)**
M. Olías, J.C. Cerón, I. Fernández, J. De la Rosa
Pollution of the aquifer with rare earth elements is documented at a site of a major spill from a mining operation.
- 65 Interaction of limestone grains and acidic solutions from the oxidation of pyrite tailings**
M. Simón, F. Martín, I. García, P. Bouza, C. Dorronsoro, J. Aguilar
Basaluminite, schwertmannite and jarosite armored the limestone grains, and almost all trace elements co-precipitated, but the precipitation of As and Pb tended to decrease as the pH rose.
- 73 Metallothionein induction, antioxidative responses, glycogen and growth changes in *Tubifex tubifex* (Oligochaete) exposed to the fungicide, fenhexamid**
Y.Y. Mosleh, S. Paris-Palacios, M. Couderchet, S. Biagianni-Risbourg, G. Vernet
*Exposure to the fungicide fenhexamid increased metallothionein levels in *Tubifex tubifex*.*
- 83 An inventory of nitric oxide emissions from soils in China**
Q. Wang, Z. Han, Y. Higano
Land use and nitrogen fertilizer use affect NO emissions from soil.
- 91 Sources of variation in concentrations of nickel and copper in mountain birch foliage near a nickel-copper smelter at Monchegorsk, north-western Russia: results of long-term monitoring**
M.V. Kozlov

Foliar concentrations of nickel and copper did not reflect emission decline during 1991–2003; annual variation was explained by weather conditions.