


Volume 140 No. 1 March 2006 ISSN 0269-7491



ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning


ASSOCIATE EDITORS

K.E. Havens

S.V. Krupa

K.C. Jones

J. W. Erisman



ENVIRONMENTAL POLLUTION

CONTENTS—Continued from outside back cover

- 79 Influence of diesel concentration on the fate of phenanthrene in soil**
A.L. Swindell, B.J. Reid
Diesel influenced the fate of [¹⁴C]
- 87 Selective supercritical fluid extraction to identify aged sediment-bound PCBs available for uptake by eel**
T. Nilsson, J. Häkkinen, P. Larsson, E. Björklund
Supercritical fluid extraction could successfully measure bioavailable fractions of PCBs in sediment.
- 95 Effluent profile of commercially used low-phosphorus fish feeds**
S.H. Sugiyura, D.D. Marchant, K. Kelsey, T. Wiggins, R.P. Ferraris
Commercial low-phosphorus fish feeds pollute less but are more costly than standard fish feeds that provide similar fish growth rates.
- 102 Release of persistent organic contaminants from carcasses of Lake Ontario Chinook salmon (*Oncorhynchus tshawytscha*)**
S. O'Toole, C. Metcalfe, I. Craine, M. Gross
Carcasses of adult Chinook salmon from Lake Ontario contribute persistent contaminants to a river ecosystem.
- 114 Comparison of natural organic acids and synthetic chelates at enhancing phytoextraction of metals from a multi-metal contaminated soil**
C.W.A. do Nascimento, D. Amarasiriwardena, B. Xing
Organic acids can be as efficient as synthetic chelates for use in phytoextraction of multi-metal contaminated soils.
- 124 Effects of inoculation of plant growth-promoting rhizobacteria on metal uptake by *Brassica juncea***
S.C. Wu, K.C. Cheung, Y.M. Luo, M.H. Wong
Rhizobacteria promoted growth above normal biomass, but did not influence plant metal concentrations.
- 136 Levels and distribution of organochlorine pesticides, polychlorinated biphenyls and polybrominated diphenyl ethers in sediments and biota from the Danube Delta, Romania**
A. Covaci, A. Gheorghe, O. Hulea, P. Schepens
Levels of persistent organohalogenated pollutants in sediments and biota from the Danube Delta are low.
- 150 Evaluating the role of desorption in bioavailability of sediment-associated contaminants using oligochaetes, semipermeable membrane devices and Tenax extraction**
M.T. Leppänen, J.V.K. Kukkonen
Desorption and animal behaviour play major roles in the availability of hydrophobic organics in sediments.
- 164 Can microbial mineralization be used to estimate microbial availability of organic contaminants in soil?**
K.T. Semple, N.M. Dew, K.J. Doick, A.H. Rhodes
Mineralization can estimate the microbial availability of ¹⁴C-contaminants in soil.
- 173 Avoidance bio-assays may help to test the ecological significance of soil pollution**
M.M. Aldaya, C. Lors, S. Salmon, J.-F. Ponge
Polluted soils are avoided by soil animals, a phenomenon which can be used as a cheap, sensitive tool for the early detection of environmental risk.
- Short communication**
- 181 Origin of 2-ethylhexanol as a VOC**
S. Nalli, O.J. Horn, A.R. Grochowalski, D.G. Cooper, J.A. Nicell
A link has been observed between the partial biodegradation of plasticizers by microorganisms and VOCs associated with poor indoor air quality.

CONTENTS

Volume 140 Number 1 2006

New initiatives

- 1 Short-term exposure to ozone altered the relative feed value of an alfalfa cultivar**
R.B. Muntifering, W.J. Manning, J.C. Lin, G.B. Robinson
Relative feed value (RFV) was affected by ozone.
- 4 Distribution and characteristics of organic micropollutants in surface sediments from Bohai Sea**
W.X. Liu, J.L. Chen, X.M. Lin, S. Tao
The recent inputs of DDT in surface sediments at some sites of Bohai Sea indicated potential ecological risk.
- 9 Aluminum-based drinking-water treatment residuals: A novel sorbent for perchlorate removal**
K.C. Makris, D. Sarkar, R. Datta
Drinking-water treatment residuals are a low-cost sorbent for perchlorate.
- 13 Uptake of vapor and particulate polycyclic aromatic hydrocarbons by cabbage**
S. Tao, X.C. Jiao, S.H. Chen, F.L. Xu, Y.J. Li, F.Z. Liu
A multivariate linear regression model was developed for predicting vegetable uptake of PAHs based on both gas and particle phases PAH concentrations.

Regular papers

- 16 Correlation of ambient inhalable bioaerosols with particulate matter and ozone: A two-year study**
A. Adhikari, T. Reponen, S.A. Grinshpun, D. Martuzevicius, G. LeMasters
Synergistic effects of these pollutants may increase incidence of respiratory health problem.
- 29 Cadmium in the Amazonian Guajará Estuary: Distribution and remobilization**
S.F. Nascimento, H. Kurzweil, W. Wruss, N. Fenzl
Distribution and remobilization of cadmium is affected differently by season.
- 43 Herbicide impact on *Hormosira banksii* gametes measured by fluorescence and germination bioassays**
C.R. Seery, L. Gunthorpe, P.J. Ralph
Chlorophyll a fluorescence measurements form the basis of a macroalgal bioassay with many advantages over germination-based methods.
- 52 Effects of heavy-metal-contaminated soil on growth, phenology and biomass turnover of *Hieracium piloselloides***
P. Ryser, W.R. Sauder
Flowering phenology shows a very sensitive response to heavy metal contamination of soils.
- 62 Seasonal variations of cadmium and zinc in *Arrhenatherum elatius*, a perennial grass species from highly contaminated soils**
A. Deram, F.-O. Denayer, D. Petit, C. Van Haluwyn
Cd and Zn bioaccumulation varies seasonally in a perennial grass.
- 71 Effects of heavy metal exposure on the condition and health of adult great tits (*Parus major*)**
T. Dauwe, E. Janssens, M. Eens
Heavy metal pollution had no clear effect on condition and health, but this may have been masked by habitat quality differences and gene flow.