

Volume 142 No. 1 July 2006 ISSN 0269-7491

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning


ASSOCIATE EDITORS

K.E. Havens

S.V. Krupa

K.C. Jones

J.W. Erisman



ENVIRONMENTAL POLLUTION

CONTENTS—Continued from outside back cover

- 83 California sea lions (*Zalophus californianus californianus*) have lower chlorinated hydrocarbon contents in northern Baja California, México, than in California, USA**
L.D. Toro, G. Heckel, V.F. Camacho-Ibar, Y. Schramm
Results suggest that sea lion prey must also have lower hydrocarbons in Baja California than in California in the USA.
- 93 Lead shot from hunting as a source of lead in human blood**
P. Johansen, H.S. Pedersen, G. Asmund, F. Riget
Birds hunted with lead shot and consumed are a source of lead in human blood.
- 98 Imposex occurrence in marine whelks at a military facility in the high Arctic**
J. Strand, C.M. Glahder, G. Asmund
Imposex as a biomarker was more sensitive for TBT than analytical chemical methods.
- 103 Bioavailability and cellular effects of metals on *Lumbricus terrestris* inhabiting volcanic soils**
A. Amaral, M. Soto, R. Cunha, I. Marigómez, A. Rodrigues
In earthworms, differences in the chloragogenous tissue morphometry may be related to the bioavailability of metals in soils.
- 109 Growth and nutritive quality of *Poa pratensis* as influenced by ozone and competition**
J. Bender, R.B. Muntifering, J.C. Lin, H.J. Weigel
*Early-season O₃ exposure decreased nutritive quality of *Poa pratensis*, and nutritive quality response to O₃ was not altered by interspecific competition.*
- 116 Effects of metals on condition and reproductive output of the smooth toadfish in Sydney estuaries, south-eastern Australia**
R. Alquezar, S.J. Markich, D.J. Booth
Enhanced metal levels in sediment reduced reproductive output and condition in toadfish.
- 123 Metal accumulation in the smooth toadfish, *Tetractenos glaber*, in estuaries around Sydney, Australia**
R. Alquezar, S.J. Markich, D.J. Booth
Metal levels in toadfish tissues reflect sediment metal levels and show gender differences.
- 132 Sorption and bioreduction of hexavalent uranium at a military facility by the Chesapeake Bay**
W. Dong, G. Xie, T.R. Miller, M.P. Franklin, T.P. Oxenberg, E.J. Bouwer, W.P. Ball, R.U. Halden
At the Aberdeen Proving Ground in Maryland, USA, migration of depleted uranium into the Chesapeake Bay is limited by rapid sorption of the radionuclide to natural organic matter followed by slow biological reduction of water-soluble U(VI) to the insoluble and less toxic U(IV) species.
- 143 Air-vegetation transfer of PCDD/PCDFs: An assessment of field data and implications for modeling**
M. Schuhmacher, K.C. Jones, J.L. Domingo
Soil and air transfer of PCDD/PCDFs to vegetation were evaluated in residential and industrial areas.
- 151 Comparison of the role of the sea club-rush *Scirpus maritimus* and the sea rush *Juncus maritimus* in terms of concentration, speciation and bioaccumulation of metals in the estuarine sediment**
C.M.R. Almeida, A.P. Mucha, M.T.S.D. Vasconcelos
*Both *Juncus maritimus* and *Scirpus maritimus* have potential for the phytostabilization of Cd, and the last one also for Pb phytostabilization.*
- 160 Fate of heavy metals after application of sewage sludge and wood-ash mixtures to short-rotation willow coppice**
I. Dimitriou, J. Eriksson, A. Adler, P. Aronsson, T. Verwijst
Commercially applied, legally regulated applications of sewage sludge and wood-ash mixtures to short-rotation willow coppice in the Enköping area, central Sweden, are tested.

CONTENTS

Volume 142 Number 1 2006

1 Urban environmental geochemistry of trace metals

C.S.C. Wong, X. Li, I. Thornton

Urban environmental geochemistry as a scientific discipline provides valuable information on trace metal contamination of the urban environment and its associated health effects.

17 Assessment of heavy metal bioavailability in contaminated sediments and soils using green fluorescent protein-based bacterial biosensors

V.H.-C. Liao, M.-T. Chien, Y.-Y. Tseng, K.-L. Ou

Nonpathogenic GFP-based bacterial biosensor is applicable in determining the bioavailability of heavy metals in environmental samples.

24 Effects of oil sands tailings compounds and harsh weather on mortality rates, growth and detoxification efforts in nestling tree swallows (*Tachycineta bicolor*)

M.-L. Gentes, C. Waldner, Z. Papp, J.E.G. Smits

Under natural stress caused by harsh weather, birds exposed to chemicals from the oil sands extraction process suffered higher mortality than those in control areas.

34 Ectomycorrhizas impede phytoremediation of polycyclic aromatic hydrocarbons (PAHs) both within and beyond the rhizosphere

E.J. Joner, C. Leyval, J.V. Colpaert

*The ectomycorrhizal fungus *S. bovinus* impeded degradation of PAHs in soil, probably due to its negative effect on the availability of mineral nutrients of more potent PAH degraders.*

39 Characterization of trace organic contaminants in marine sediment from Yeongil Bay, Korea: 1. Instrumental analyses

C.-H. Koh, J.S. Khim, D.L. Villeneuve, K. Kannan, J.P. Giesy

Among various sediment contaminant classes measured, nonylphenol and PAHs are responsible for the variability among sampling sites, suggesting the existence of multiple sources in Yeongil Bay sediment.

48 Characterization of trace organic contaminants in marine sediment from Yeongil Bay, Korea: 2. Dioxin-like and estrogenic activities

C.-H. Koh, J.S. Khim, D.L. Villeneuve, K. Kannan, J.P. Giesy

In vitro bioassay responses observed for Yeongil Bay surficial sediment and sediment core extracts showed the greatest dioxin-like and estrogenic activities in the mid-polarity fraction containing PAHs as well as chlorinated dioxins and furans.

58 Effects of NO₂ and NH₃ from road traffic on epiphytic lichens

L. Frati, E. Caprasecca, S. Santoni, C. Gaggi, A. Guttova, S. Gaudino, A. Pati, S. Rosamilia, S.A. Pirintsos, S. Loppi

NH₃ and NO₂ from a busy road in central Italy do not influence epiphytic lichens.

65 Genotoxicity biomarkers and acetylcholinesterase activity in natural populations of *Mytilus galloprovincialis* along a pollution gradient in the Gulf of Oristano (Sardinia, western Mediterranean)

P. Magni, G. De Falco, C. Falugi, M. Franzoni, M. Monteverde, E. Perrone, M. Sgro, C. Bolognesi

Biomarkers can be useful in coastal marine biomonitoring.

73 *Brachidontes variabilis* and *Patella* sp. as quantitative biological indicators for cadmium, lead and mercury in the Lebanese coastal waters

K.F. Nakhlé, D. Cossa, G. Khalaf, B. Beliaeff

Two molluscs were effective bioindicators for metal pollution in waters along the Lebanese coast.