


Volume 134 No. 2 March 2005 ISSN 0269-7491



ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning


ASSOCIATE EDITORS

K.E. Havens

S.V. Krupa

K.C. Jones

J. W. Erisman



ENVIRONMENTAL POLLUTION

CONTENTS—Continued from outside back cover

- 257 **Interactive effect of *Brevibacillus brevis* and *Glomus mosseae*, both isolated from Cd contaminated soil, on plant growth, physiological mycorrhizal fungal characteristics and soil enzymatic activities in Cd polluted soil**
A. Vivas, J.M. Barea, R. Azcón
Endemic isolates of bacteria and fungi were effective in bioremediation.
- 267 **Oil spill in the Río de la Plata estuary, Argentina: 2-hydrocarbon disappearance rates in sediments and soils**
J.C. Colombo, A. Barreda, C. Bilos, N. Cappelletti, M.C. Migoya, C. Skorupka
Hydrocarbon disappearance rates in Río de la Plata sediments and soils were determined.
- 277 **Oil spill in the Río de la Plata estuary, Argentina: 1. Biogeochemical assessment of waters, sediments, soils and biota**
J.C. Colombo, A. Barreda, C. Bilos, N. Cappelletti, S. Demichelis, P. Lombardi, M.C. Migoya, C. Skorupka, G. Suárez
The Oil spill impact in the Río de la Plata was evaluated by HRGC and multivariate techniques applied to waters, sediments and biota.
- 291 **Ozone exposure induces the activation of leaf senescence-related processes and morphological and growth changes in seedlings of Mediterranean tree species**
A. Ribas, J. Peñuelas, S. Elvira, B.S. Gimeno
Ozone induces species-specific leaf senescence-related process and morphological and growth changes in seedlings of Mediterranean tree species.
- 301 **Review of disposal technologies for chromated copper arsenate (CCA) treated wood waste, with detailed analyses of thermochemical conversion processes**
L. Helsen, E. Van den Bulck
Low-temperature pyrolysis or high-temperature gasification are the best recyclable solutions.
- 315 **PAH dissipation in a contaminated river sediment under oxic and anoxic conditions**
C. Quantin, E.J. Joner, J.M. Portal, J. Berthelin
Natural attenuation of PAHs under anaerobic conditions is exceedingly slow.
- 323 **PCBs and DDT in the serum of juvenile California sea lions: associations with vitamins A and E and thyroid hormones**
C. Debier, G.M. Ylitalo, M. Weise, F. Gulland, D.P. Costa, B.J. Le Boeuf, T. de Tillesse, Y. Larondelle
Results show high levels of organochlorine contaminants in juvenile California sea lions and a link between vitamin A, thyroid hormones and PCB exposure.
- 333 **A lysimeter experiment to investigate the leaching of veterinary antibiotics through a clay soil and comparison with field data**
P. Kay, P.A. Blackwell, A.B.A. Boxall
This paper describes one of the first studies to investigate the fate of veterinary medicines in cracking clay soils.
- 343 **Screening the wetland plant species *Alisma plantago-aquatica*, *Carex rostrata* and *Phalaris arundinacea* for innate tolerance to zinc and comparison with *Eriophorum angustifolium* and *Festuca rubra* Merlin**
D.J. Matthews, B.M. Moran, M.L. Otte
Some wetland angiosperms are tolerant to high concentrations of metals, regardless of conditions in the plants' natural habitat.
- 353 **Assessing breeding potential of peregrine falcons based on chlorinated hydrocarbon concentrations in prey**
J.E. Elliott, M.J. Miller, L.K. Wilson
Based on the level of DDE contamination of prey items, it seems unlikely that peregrine falcons could breed successfully throughout most of the Okanagan Valley of British Columbia.

CONTENTS

Volume 134 Number 2 2005

Research papers

- 181 Assessing soil ecotoxicity of methyl tert-butyl ether using earthworm bioassay; closed soil microcosm test for volatile organic compounds**
Y.-J. An
Earthworm bioassay can be a good protocol to assess soil ecotoxicity of volatile organic compounds such as MTBE.
- 187 Effect of organic fertilizers derived dissolved organic matter on pesticide sorption and leaching**
K. Li, B. Xing, W.A. Torello
Dissolved organic matter could result in enhanced transport of chemicals applied to turf.
- 195 Microfibril angle in wood of Scots pine trees (*Pinus sylvestris*) after irradiation from the Chernobyl nuclear reactor accident**
M. Tulik, A. Rusin
The angle of microfibrils in the S₂ layer in wood deposited after the Chernobyl incident was changed.
- 203 Absence of Hg transpiration by shoot after Hg uptake by roots of six terrestrial plant species**
M. Greger, Y. Wang, C. Neuschütz
Mercury translocation to shoots was low.
- 209 Effects of elevated ozone on leaf $\delta^{13}\text{C}$ and leaf conductance of plant species grown in semi-natural grassland with or without irrigation**
M. Jäggi, M. Saurer, M. Volk, J. Fuhrer
Under natural field conditions, lack of precipitation may not protect semi-natural vegetation from O₃ effects on leaf gas exchange.
- 217 Evidence of the involvement of plant ligno-cellulosic structure in the sequestration of Pb: an X-ray spectroscopy-based analysis**
M. Marmiroli, G. Antonioli, E. Maestri, N. Marmiroli
*Lignin and cellulose can play a major role in the accumulation of lead in *Juglans regia* roots.*
- 229 Nitrogen species in drinking water indicate potential exposure pathway for Balkan Endemic Nephropathy**
N. Niagolova, S.P. McElmurry, T.C. Voice, D.T. Long, E.A. Petropoulos, I. Havezov, K. Chou, V. Ganey
Water in villages affected and unaffected by Balkan Endemic Nephropathy had no significant differences in nitrogen compounds.
- 239 Cadmium uptake by tobacco as affected by liming, N form, and year of cultivation**
C.D. Tsadilas, N.A. Karaivazoglou, N.C. Tsotsolis, S. Stamatiadis, V. Samaras
Liming and N form affect Cd uptake by Virginia tobacco which contributes significantly to the great reduction of extractable soil Cd after three years of continuous cultivation.
- 247 Isotopic characterisation of lead in contaminated soils from the vicinity of a non-ferrous metal smelter near Plovdiv, Bulgaria**
J.R. Bacon, N.S. Dinev
Although soils in the vicinity of a non-ferrous metal smelter near Plovdiv, Bulgaria, have become highly contaminated with the ores used, lead isotope analysis has revealed that up to 12% of current deposition could be from other sources such petrol lead.