

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa

K.C. Jones J. W. Erisman

ENVIRONMENTAL POLLUTION

CONTENTS - Continued from outside back cover

293 The Solling roof revisited – slow recovery from acidification observed and modeled despite a decade of "clean-rain" treatment

L. Martinson, N. Lamersdorf, P. Warfvinge

Model predictions show slow recovery by 2050.

The effect of hydrological regime on the metal bioavailability for the wetland plant species Salix cinerea B. Vandecasteele, P. Quataert, F.M.G. Tack

A hydrological regime aiming at wetland creation is a potential management option that favors reducing Cd plant availability in polluted freshwater wetlands.

313 Manganese toxicity thresholds for restoration grass species M.W. Paschke, A. Valdecantos, E.F. Redente

Mn phytotoxicity thresholds for restoration grasses should be useful for risk assessments of metal-contaminated lands.

323 Decomposition of leaves of the metallophyte *Arabidopsis halleri* in soil microcosms: fate of Zn and Cd from plant residues

U. Boucher, I. Lamy, P. Cambier, M. Balabane

Changes in metallic speciation occurred in two stages.

333 Effects of arsenic on concentration and distribution of nutrients in the fronds of the arsenic hyperaccumulator Pteris vittata L.

C. Tu, L.Q. Ma

Arsenic hyperaccumulator P. vittata maintained adequate levels of essential nutrients in the fronds under arsenic stress.

Letter to the editor

Comment on "Characterization of a reference site for quantifying uncertainties related to soil sampling" by S. Barbizzi et al. (2004)

P. Baveye



ENVIRONMENTAL POLLUTION

www.elsevier.com/locate/envpol

CONTENTS

Volume 135 Number 2 2005

187 Decomposition in soil microcosms of leaves of the metallophyte *Arabidopsis halleri*: effect of leaf-associated heavy metals on biodegradation

U. Boucher, M. Balabane, I. Lamy, P. Cambier

Decomposition of slowly decomposable fractions of metal-rich plant residues is slowed by the metals.

195 Sub-cellular partitioning of Cd, Cu and Zn in tissues of indigenous unionid bivalves living along a metal exposure gradient and links to metal-induced effects

E. Bonneris, O. Perceval, S. Masson, L. Hare, P.G.C. Campbell

In natural populations of bivalves, metals were differentially bound in the gills and digestive gland.

209 Metal stress consequences on frost hardiness of plants at northern high latitudes: a review and hypothesis K. Taulavuori, M.N.V. Prasad, E. Taulavuori, K. Laine

Metal stress may reduce plant frost hardiness.

221 Dynamical coupling of PBPK/PD and AUC-based toxicity models for arsenic in tilapia *Oreochromis mossambicus* from blackfoot disease area in Taiwan

C.-M. Liao, H.-M. Liang, B.-C. Chen, S. Singh, J.-W. Tsai, Y.-H. Chou, W.-T. Lin

Integrated toxicity models can identify dynamic responses of fish to arsenic.

235 Octanol-solubility of dissolved and particulate trace metals in contaminated rivers: implications for metal reactivity and availability

A. Turner, E. Mawji

New approaches are presented for fractionating trace metals in natural waters.

245 Dietary mercury exposure and bioaccumulation in amphibian larvae inhabiting Carolina bay wetlands J.M. Unrine, C.H. Jagoe, A.C. Brinton, H.A. Brant, N.T. Garvin

Tadpoles accumulate significant concentrations of mercury in isolated wetlands.

255 Diversity of arbuscular mycorrhizal fungi in grassland spontaneously developed on area polluted by a fertilizer plant C. Renker, V. Blanke, F. Buscot

Almost all plants species were mycorrhizal.

267 In vitro toxicity of selected pesticides on RTG-2 and RTL-W1 fish cell lines M.M. Babín, J.V. Tarazona

In vitro cell cultures can provide sensitive indicators for pesticide effects on biota.

275 Early-phase immunodetection of metallothionein and heat shock proteins in extruded earthworm coelomocytes after dermal exposure to metal ions

J. Homa, E. Olchawa, S.R. Stürzenbaum, A. John Morgan, B. Plytycz

Metals upregulate stress response proteins in earthworm coelomocytes.

- 281 Oxygen levels versus chemical pollutants: do they have similar influence on macrofaunal assemblages? A case study in a harbour with two opposing entrances

 J.M. Guerra-García, J.C. García-Gómez
 - A high biomass of benthic animals was found in a metal-polluted estuary where high oxygen levels occurred.