


Volume 140 No. 2 March 2006 ISSN 0269-7491



ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning


ASSOCIATE EDITORS

K.E. Havens

S.V. Krupa

K.C. Jones

J. W. Erisman



ENVIRONMENTAL POLLUTION

CONTENTS—Continued from outside back cover

- 304 Baseline mercury and zinc concentrations in terrestrial and coastal organisms of Admiralty Bay, Antarctica**
I.R. dos Santos, E.V. Silva-Filho, C. Schaefer, S. Maria Sella, C.A. Silva, V. Gomes, M.J.d.A.C.R. Passos, P. Van Ngan
Low levels of mercury and zinc occurred in soil and plant samples from Antarctica, but high levels occurred in birds and mammals.
- 312 Organochlorine contaminant and retinoid levels in blubber of common dolphins (*Delphinus delphis*) off northwestern Spain**
V. Tornero, A. Borrell, A. Aguilar, J. Forcada, C. Lockyer
Organochlorine contaminants and retinoids in common dolphins.
- 322 Effect of the cationic composition of sorption solution on the quantification of sorption-desorption parameters of heavy metals in soils**
J. Sastre, G. Rauret, M. Vidal
Sorption solution composition modifies metal sorption-desorption pattern in soils.
- 340 Effect of humic acid on water chemistry, bioavailability and toxicity of aluminium in the freshwater snail, *Lymnaea stagnalis*, at neutral pH**
A. Dobranskyte, R. Jugdaohsingh, C.R. McCrohan, E. Stuchlik, J.J. Powell, K.N. White
Humic acid reduces aluminium toxicity in freshwater snails.
- 348 Pesticide residues in Nile tilapia (*Oreochromis niloticus*) and Nile perch (*Lates niloticus*) from Southern Lake Victoria, Tanzania**
L. Henry, M.A. Kishimba
Fish from Lake Victoria had relatively low pesticide levels.
- 355 Spatio-temporal and species-specific variation in PBDE levels/patterns in British Columbia's coastal waters**
M.G. Ikonomou, M.P. Fernandez, Z.L. Hickman
PBDEs in biota and sediments from the West Coast of Canada reflect temporal, species-specific and geographic distributions.
- 364 The use of directional traps for the assessment of short-term phenanthrene effects upon soil springtail communities**
F. Gass, S. Gillet, J.-F. Ponge
Soil springtails may be attracted to phenanthrene in dosing trials.
- 371 Occurrence of organochlorine pesticides (OCPs) and their enantiomeric signatures, and concentrations of polybrominated diphenyl ethers (PBDEs) in the Adélie penguin food web, Antarctica**
S. Corsolini, A. Covaci, N. Ademollo, S. Focardi, P. Schepens
Enantiomeric signature and accumulation of polybrominated diphenyl ethers in the Adélie Penguin food chain are discussed.

CONTENTS

Volume 140 Number 2 2006

- 187 Nutrient status and plant growth effects of forest soils in the Basin of Mexico**
M.E. Fenn, V.M. Perea-Estrada, L.I. de Bauer, M. Pérez-Suárez, D.R. Parker, V.M. Cetina-Alcalá
Plant response to N deposition may be limited by P limitation in forests growing on Andisol soils in the Basin of Mexico.
- 200 Development of the high spatial resolution EMICAT2000 emission model for air pollutants from the north-eastern Iberian Peninsula (Catalonia, Spain)**
R. Parra, P. Jiménez, J.M. Baldasano
A high spatial resolution and multipurpose emission model is presented.
- 220 Preliminary assessment of risk of ozone impacts to maize (*Zea mays*) in southern Africa**
A.M. van Tienhoven, M. Zunckel, L. Emberson, A. Koosailee, L. Otter
Ozone impacts on crops in southern Africa are assessed using current European-based methods (AOT40 concept).
- 231 Development and application of a sediment toxicity test using the benthic cladoceran *Chydorus sphaericus***
T. Dekker, G.D. Greve, T.L. Ter Laak, M.E. Boivin, B. Veuger, G. Gortzak, S. Dumfries, S.M.G. Lücker, M.H.S. Kraak, W. Admiraal, H.G. van der Geest
A test was developed for assaying sediment toxicity using a commonly occurring small-bodied cladoceran.
- 239 Functional recovery of biofilm bacterial communities after copper exposure**
M.-E.Y. Boivin, B. Massieux, A.M. Breure, G.D. Greve, M. Rutgers, W. Admiraal
Genetically different bacterial communities can have similar functions and tolerance to copper.
- 247 Effects of road de-icing salt (NaCl) on larval wood frogs (*Rana sylvatica*)**
D. Sanzo, S.J. Hecnar
Road salts have toxic effects on amphibians at environmentally realistic concentrations.
- 257 Sources of errors in assessing ozone visible symptoms on native vegetation**
F. Bussotti, M. Schaub, A. Cozzi, G. Gerosa, K. Novak, C. Hug
The quality of foliar ozone-symptom assessment depends on the experience of the surveyor, the symptom specificity, and the symptom variability within the same species.
- 269 Miniaturized test system for soil respiration induced by volatile pollutants**
K. Kaufmann, S.J. Chapman, C.D. Campbell, H. Harms, P. Höhener
A new method to study soil respiration is used when volatile organic contaminants are added.
- 279 Assessment of the bioaccessibility of polycyclic aromatic hydrocarbons in soils from Beijing using an in vitro test**
X.-Y. Tang, L. Tang, Y.-G. Zhu, B.-S. Xing, J. Duan, M.-H. Zheng
Oral bioaccessibility of soil total PAHs in small intestinal condition was significantly higher than gastric condition.
- 286 The effect of emergent macrophytes on the dynamics of sulfur species and trace metals in wetland sediments**
J.H. Choi, S.S. Park, P.R. Jaffé
Plants affected the biogeochemistry of sulfur species and the mobility of trace metals in wetland sediments.
- 294 Methyl tert-butyl ether (MTBE) in finished drinking water in Germany**
A. Kolb, W. Püttmann
MTBE in drinking water comes from different sources of MTBE in the aquatic environment.