


Volume 141 No. 2 May 2006 ISSN 0269-7491



ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning


ASSOCIATE EDITORS

K.E. Havens

S.V. Krupa

K.C. Jones

J.W. Erisman



ENVIRONMENTAL POLLUTION

CONTENTS—Continued from outside back cover

- 275 **Ozone sensitivity of currant tomato (*Lycopersicon pimpinellifolium*), a potential bioindicator species**
M. Iriti, L. Belli, C. Nali, G. Lorenzini, G. Gerosa, F. Faoro
*Currant tomato (*Lycopersicon pimpinellifolium*) was found to be very sensitive to ozone.*
- 283 **Ecological impact assessment of metallurgic effluents using in situ biomarker assays**
W. De Coen, J. Robbens, C. Janssen
In situ monitoring of metallurgic effluents using biomarker analysis.
- 295 **Natural and anthropogenic geochemical signatures of floodplain and deltaic sedimentary strata, Sacramento-San Joaquin Delta, California, USA**
G.B. Pasternack, K.J. Brown
Geomorphic-geochemical linkages on an upper deltaic plain have dynamics that defy expectations for achieving restoration "success".
- 310 **Influence of hydrology on heavy metal speciation and mobility in a Pb-Zn mine tailing**
E. Kovács, W.E. Dubbin, J. Tamás
Variable hydrology influences heavy metal speciation and mobility, and the formation of neutralization zones, in a Pb-Zn mine tailing.
- 321 **Antimony mobility in Japanese agricultural soils and the factors affecting antimony sorption behavior**
Y. Nakamaru, K. Tagami, S. Uchida
Antimony mobility in tested Japanese agricultural soils was generally low and was affected by pH and phosphate concentration.
- 327 **Polychlorinated biphenyls (PCBs) in soils of the Moscow region: Concentrations and small-scale distribution along an urban-rural transect**
W. Wilcke, M. Krauss, G. Safronov, A.D. Fokin, M. Kaupenjohann
PCBs are accumulated on aggregate surfaces where soil biological activity is particularly high.
- 336 **Trace metals in sediments of two estuarine lagoons from Puerto Rico**
D. Acevedo-Figueroa, B.D. Jiménez, C.J. Rodríguez-Sierra
- 343 **Multigenerational cadmium acclimation and biokinetics in *Daphnia magna***
R. Guan, W.-X. Wang
*Tolerance of sensitive *Daphnia magna* clones to cadmium was dependent on long-term or multigenerational exposure.*
- 353 **Effect of chromium on the fatty acid composition of two strains of *Euglena gracilis***
I. Rocchetta, M. Mazzuca, V. Conforti, L. Ruiz, V. Balzaretto, M.d.C.R. de Molina
*Fatty acid evaluation in the presence of chromium in *Euglena gracilis* grown in different culture conditions.*
- 359 **Remobilization of metals from slag and polluted sediments (Case Study: The canal of the Deûle River, northern France)**
N. Vdović, G. Billon, C. Gabelle, J.-L. Potdevin
Heavily polluted sediments of the Deûle River are still able to adsorb heavy metals released from toxic wastes.
- 370 **Relation between PAH and black carbon contents in size fractions of Norwegian harbor sediments**
A.M.P. Oen, G. Cornelissen, G.D. Breedveld
PAH contents correlated better with black carbon than organic carbon for four Norwegian harbor sediments.
- 381 **High average daily intake of PCDD/Fs and serum levels in residents living near a deserted factory producing pentachlorophenol (PCP) in Taiwan: Influence of contaminated fish consumption**
C.C. Lee, W.T. Lin, P.C. Liao, H.J. Su, H.L. Chen
Inhabitants living near a deserted PCP factory are exposed to high PCDD/F levels.

CONTENTS

Volume 141 Number 2 2006

- 195 Removal of cyanobacterial blooms in Taihu Lake using local soils. I. Equilibrium and kinetic screening on the flocculation of *Microcystis aeruginosa* using commercially available clays and minerals**
G. Pan, M.-M. Zhang, H. Chen, H. Zou, H. Yan
Sepiolite was the most effective flocculant among 26 commercially available clays and minerals in removing harmful algal cells from freshwaters.
- 201 Removal of cyanobacterial blooms in Taihu Lake using local soils. II. Effective removal of *Microcystis aeruginosa* using local soils and sediments modified by chitosan**
H. Zou, G. Pan, H. Chen, X. Yuan
Chitosan modification can turn many solids, such as local clays and soils, into highly effective flocculants in removing harmful cyanobacterial blooms in freshwaters.
- 206 Removal of harmful cyanobacterial blooms in Taihu Lake using local soils. III. Factors affecting the removal efficiency and an in situ field experiment using chitosan-modified local soils**
G. Pan, H. Zou, H. Chen, X. Yuan
A loading of 0.025 g/L chitosan-modified local soils removed 99% algal cells in 16 h in a field enclosure of Taihu Lake.
- 213 Polychlorinated dibenzo-*p*-dioxins, dibenzofurans, biphenyls, naphthalenes and polybrominated diphenyl ethers in the edible fish caught from the Baltic Sea and lakes in Finland**
P. Isosaari, A. Hallikainen, H. Kiviranta, P.J. Vuorinen, R. Parmanne, J. Koistinen, T. Vartiainen
Lake fish is less contaminated with dioxins, PCBs and chloronaphthalenes than Baltic Sea fish but similarly contaminated with polybrominated diphenyl ethers.
- 226 Temporal trend of bis(4-chlorophenyl) sulfone, methylsulfonyl-DDE and -PCBs in Baltic guillemot (*Uria aalge*) egg 1971–2001 – A comparison to 4,4'-DDE and PCB trends**
H. Jörundsdóttir, K. Norström, M. Olsson, H. Pham-Tuan, H. Hühnerfuss, A. Bignert, Å. Bergman
The temporal trend of the plastic monomer bis(4-chlorophenyl) sulfone (BCPS) showed that it was present in low ppm concentrations in Baltic guillemot eggs and decreased by only 1.6% annually, 1971–2001.
- 238 Arsenic speciation, and arsenic and phosphate distribution in arsenic hyperaccumulator *Pteris vittata* L. and non-hyperaccumulator *Pteris ensiformis* L.**
N. Singh, L.Q. Ma
Pteris vittata may effectively reduce arsenite in its fronds.
- 247 Environmental evidence of fossil fuel pollution in Laguna Chica de San Pedro lake sediments (Central Chile)**
L. Chirinos, N.L. Rose, R. Urrutia, P. Muñoz, F. Torrejón, L. Torres, F. Cruces, A. Araneda, C. Zaror
The lake sediment record of SCPs shows the record of fossil-fuel derived pollution in Central Chile.
- 257 Characterizing the risk assessment of heavy metals and sampling uncertainty analysis in paddy field by geostatistics and GIS**
X. Liu, J. Wu, J. Xu
Probability maps gave insight into risk assesment of environmental metals in a rice paddy field.
- 265 Persistent organic pollutants and sedimentary organic matter properties: A case study in the Kishon River, Israel**
A. Oren, Z. Aizenshtat, B. Chefetz
Petrogenic-derived xenobiotics affect the properties of sedimentary organic matter and its sorption capabilities.