


Volume 142 No. 2 July 2006 ISSN 0269-7491



# ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

**W.J. Manning**


ASSOCIATE EDITORS

**K.E. Havens**

**S.V. Krupa**

**K.C. Jones**

**J.W. Erisman**



# ENVIRONMENTAL POLLUTION

## CONTENTS—Continued from outside back cover

- 288 **Influence of vegetation in mitigation of methyl parathion runoff**  
M.T. Moore, E.R. Bennett, C.M. Cooper, S. Smith Jr., J.L. Farris, K.G. Drouillard, R. Schulz  
*Vegetated wetlands are three times more effective at mitigating methyl parathion runoff than are non-vegetated wetlands.*
- 295 **Evaluating the vulnerability of surface waters to antibiotic contamination from varying wastewater treatment plant discharges**  
A.L. Batt, I.B. Bruce, D.S. Aga  
*This work investigates the extent of antibiotic concentrations in receiving waters from discharges of wastewater treatment plants.*
- 303 **Metal stoichiometry in predicting Cd and Cu toxicity to a freshwater green alga *Chlamydomonas reinhardtii***  
W.-X. Wang, R.C.H. Dei  
*The stoichiometry of metals and P can be used to predict metal toxicity to freshwater phytoplankton under different P conditions.*
- 313 **Seasonal variations in hepatic Cd and Cu concentrations and in the sub-cellular distribution of these metals in juvenile yellow perch (*Perca flavescens*)**  
L.D. Kraemer, P.G.C. Campbell, L. Hare  
*In fish from metal-contaminated sites, seasonal variations in hepatic Cd and Cu concentrations were greater than in fish from reference sites, and homeostatic control of sub-cellular metal distribution was compromised.*
- 326 **Electrokinetic enhancement of phenanthrene biodegradation in creosote-polluted clay soil**  
J.-L. Niqui-Arroyo, M. Bueno-Montes, R. Posada-Baquero, J.-J. Ortega-Calvo  
*Electrokinetic bioremediation is a potentially effective technology to treat PAH-polluted, clay-rich soils.*
- 333 **Calibration of the Chemcatcher passive sampler for the monitoring of priority organic pollutants in water**  
B. Vrana, G.A. Mills, E. Dominiak, R. Greenwood  
*A calibration method and data that are required for in situ measurement of the time-weighted average concentrations of hydrophobic priority organic pollutants in water using Chemcatcher passive sampling device are presented.*
- 344 **Herbicide contamination of surficial groundwater in Northern Italy**  
L. Guzzella, F. Pozzoni, G. Giuliano  
*Herbicide groundwater pollution is an environmental problem in Northern Italy.*
- 354 **Field responses of *Prunus serotina* and *Asclepias syriaca* to ozone around southern Lake Michigan**  
J.P. Bennett, E.A. Jepsen, J.A. Roth  
*Decreased cherry branch elongation, milkweed stem height and pod production, and foliar injury on both species occurred at sites around southern Lake Michigan at ozone exposures of 13 SUM06 ppm-h and 93–98 ppb peak hourly.*
- 367 **Perfluorooctanesulfonate and related fluorochemicals in biological samples from the north coast of Colombia**  
J. Olivero-Verbel, L. Tao, B. Johnson-Restrepo, J. Guette-Fernández, R. Baldiris-Avila, I. O'byrne-Hoyos, K. Kannan  
*Perfluorooctanesulfonate and related perfluorinated compounds have been found in a tropical ecosystem of South America.*
- 373 **Heavy metals affect the coelomocyte-bacteria balance in earthworms: Environmental interactions between abiotic and biotic stressors**  
E. Olchawa, M. Bzowska, S.R. Stürzenbaum, A.J. Morgan, B. Plytycz  
*Metals affect coelomocyte-bacterial balance in earthworms.*

**CONTENTS**

**Volume 142 Number 2 2006**

- 191 Biodegradation and speciation of residual SS-ethylenediaminedisuccinic acid (EDDS) in soil solution left after soil washing**  
S. Tandy, A. Ammann, R. Schulin, B. Nowack  
*Even in polluted soils, EDDS is degraded.*
- 200 Use of hydra for chronic toxicity assessment of waters intended for human consumption**  
V.V. Arkhipchuk, C. Blaise, M.V. Malinovskaya  
*Hydra is a reliable and relevant test organism for the assessment of acute and chronic toxicity of waters intended for human consumption.*
- 212 Consequences for *Protaphorura armata* (Collembola: Onychiuridae) following exposure to genetically modified *Bacillus thuringiensis* (Bt) maize and non-Bt maize**  
L.-H. Heckmann, B.S. Griffiths, S. Caul, J. Thompson, M. Pusztai-Carey, W.J. Moar, M.N. Andersen, P.H. Krogh  
*Protaphorura armata performed equally well when reared on two Bt and three non-Bt maize varieties.*
- 217 Physicochemical characterization and mercury speciation of particle-size soil fractions from an abandoned mining area in Mieres, Asturias (Spain)**  
R. Fernández-Martínez, J. Loredó, A. Ordóñez, M.I. Rucandío  
*Factors affecting mercury mobility in soils influenced by abandoned mining.*
- 227 A spatially-evaluated methodology for assessing risk to a population from contaminated land**  
J.R. Gay, A. Korre  
*A methodology for calculating and mapping risks to a population from intake of soil contaminants.*
- 235 Polychlorinated biphenyl congener patterns in tree swallows (*Tachycineta bicolor*) nesting in the Housatonic River watershed, western Massachusetts, USA, using a novel statistical approach**  
C.M. Custer, L.B. Read  
*Polychlorinated biphenyl congener patterns in swallows differed between the main stem of the Housatonic River, MA and other locations in the watershed.*
- 246 Impact of short-term and long-term elevated CO<sub>2</sub> on emission of carbonyls from adult *Quercus petraea* and *Carpinus betulus* trees**  
J. Kreuzwieser, H. Rennenberg, R. Steinbrecher  
*Carbonyl exchange of European tree species was not affected by elevated ambient carbon dioxide concentrations.*
- 254 Spatial variability and temporal stability of throughfall deposition under beech (*Fagus sylvatica* L.) in relationship to canopy structure**  
J. Staelens, A. De Schrijver, K. Verheyen, N.E.C. Verhoest  
*Ion deposition under a deciduous beech tree was strongly affected by the canopy structure throughout the entire year.*
- 264 The marine macroalga *Cystoseira baccata* as biosorbent for cadmium(II) and lead(II) removal: Kinetic and equilibrium studies**  
P. Lodeiro, J.L. Barriada, R. Herrero, M.E. Sastre de Vicente  
*Marine macroalgae show promise for biosorption of lead and cadmium.*
- 274 Benthic foraminifera and heavy metals distribution: A case study from the Naples Harbour (Tyrrhenian Sea, Southern Italy)**  
L. Ferraro, M. Sprovieri, I. Alberico, F. Lirer, L. Prevedello, E. Marsella  
*Benthic foraminiferal density and species distribution may be used as pollution indicators.*