


Volume 133 No. 3 February 2005 ISSN 0269-7491



# ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

**W.J. Manning**


ASSOCIATE EDITORS

**K.E. Havens**

**S.V. Krupa**

**K.C. Jones**

**J. W. Erisman**





# ENVIRONMENTAL POLLUTION

## CONTENTS—Continued from outside back cover

- 489 **Accumulation of butyl- and phenyltin compounds in starfish and bivalves from the coastal environment of Korea**  
W.J. Shim, U.H. Yim, N.S. Kim, S.H. Hong, J.R. Oh, J.K. Jeon, H. Okamura  
*Starfish are effective organisms for monitoring phenyltin contamination in the marine environment.*
- 501 **Brominated flame retardants in *Alburnus alburnus* from Cinca River Basin (Spain)**  
E. Eljarrat, A. de la Cal, D. Raldua, C. Duran, D. Barcelo  
*PBDEs and hexabromocyclododecane (HBCD) are shown to accumulate in fish and data are presented upstream and around an industrial area.*
- 509 **Temporal trends of organochlorine contamination in Black Guillemots in Iceland from 1976 to 1996**  
K. Ólafsdóttir, Æ. Petersen, E.V. Magnúsdóttir, T. Björnsson, T. Jóhannesson  
*Baseline trends for a range of organochlorine pesticides in the marine environment of the North Atlantic can be inferred.*
- 517 **Evaluation of photolysis and hydrolysis of atrazine and its first degradation products in the presence of humic acids**  
H. Prosen, L. Zupančič-Kralj  
*Different humic acid-influenced degradation processes influence atrazine and its degradation products.*
- 531 **Differential swimming performance of two natricine snakes exposed to a cholinesterase-inhibiting pesticide**  
W.A. Hopkins, C.T. Winne, S.E. DuRant  
*Exposure to a cholinesterase inhibitor reduces swimming velocity in snakes.*
- 541 **Clonal variation in heavy metal accumulation and biomass production in a poplar coppice culture. II. Vertical distribution and phytoextraction potential**  
I. Laureysens, L. De Temmerman, T. Hastir, M. Van Gysel, R. Ceulemans  
*Poplar shows potential for phytoextraction of Al, Cd and Zn on slightly contaminated soils.*
- 553 **Physiological and foliar symptom response in the crowns of *Prunus serotina*, *Fraxinus americana* and *Acer rubrum* canopy trees to ambient ozone under forest conditions**  
M. Schaub, J.M. Skelly, J.W. Zhang, J.A. Ferdinand, J.E. Savage, R.E. Stevenson, D.D. Davis, K.C. Steiner  
*Within the heterogeneous environment of a mature forest, many factors in addition to soil moisture play a significant role in determining exposure/response relationships to ozone.*
- 569 **Stochastic analysis to assess the spatial distribution of groundwater nitrate concentrations in the Po catchment (Italy)**  
S. Cinnirella, G. Buttafuoco, N. Pirrone  
*The stochastic simulation should be preferred to kriging in environmental studies, whenever it is critical to preserve the variation of a variable.*
- 581 **Absorption of decabromodiphenyl ether and other organohalogen chemicals by grey seals (*Halichoerus grypus*)**  
G.O. Thomas, S.E.W. Moss, L. Asplund, A.J. Hall  
*Decabromodiphenyl ether is absorbed effectively from the diet by grey seals, and can be stored in the blubber even after exposure ceases.*
- 587 **Spray irrigation of landfill leachate: estimating potential exposures to workers and bystanders using a modified air box model and generalised source term**  
D. Gray, S.J.T. Pollard, L. Spence, R. Smith, J.R. Gronow  
*Modelling approaches are used to assess human exposure routes to chemicals during spray irrigation to landfill leachates.*



**CONTENTS**

**Volume 133 Number 3 2005**

- 389 Aquatic microcosm assessment of the effects of tylosin on *Lemna gibba* and *Myriophyllum spicatum***  
R.A. Brain, K.(Jim) Bestari, H. Sanderson, M.L. Hanson, C.J. Wilson, D.J. Johnson, P.K. Sibley, K.R. Solomon  
*Tylosin is not expected to have ecologically significant effects on Ontario freshwater macrophytes.*
- 403 Monitoring and modelling of biosphere/atmosphere exchange of gases and aerosols in Europe**  
J.W. Erisman, A. Vermeulen, A. Hensen, C. Flechard, U. Dämmgen, D. Fowler, M. Sutton, L. Grünhage, J.-P. Tuovinen  
*Monitoring and modelling of the deposition of sulphur and nitrogen components and the exposure of ozone has gained much progress through the research within BIATEX.*
- 415 Concentrations and compositions of organochlorine contaminants in sediments, soils, crustaceans, fishes and birds collected from Lake Tai, Hangzhou Bay and Shanghai city region, China**  
H. Nakata, Y. Hirakawa, M. Kawazoe, T. Nakabo, K. Arizono, S.-I. Abe, T. Kitano, H. Shimada, I. Watanabe, W. Li, X. Ding  
*Elevated concentrations of DDTs were detected in sediments, soils, and wildlife collected from China.*
- 431 Modelling the fate of sulphur-35 in crops. 1. Calibration data**  
C. Collins, N. Cunningham  
*The deposition, fate and loss of <sup>35</sup>S in crops were quantified following exposure to COS.*
- 439 Modelling the fate of sulphur-35 in crops. 2. Development and validation of the CROPS-35 model**  
C. Collins, N. Cunningham  
*The calibration and validation of a model for the prediction of the fate of <sup>35</sup>S in vegetation is described.*
- 447 Intrinsic and induced isoproturon catabolic activity in dissimilar soils and soils under dissimilar land use**  
B.J. Reid, N.D. Papanikolaou, R.K. Wilcox  
*Dissimilar levels of isoproturon catabolic activity in dissimilar soils and soils under dissimilar land use influence inferred risk.*
- 455 The effects of perennial ryegrass and alfalfa on microbial abundance and diversity in petroleum contaminated soil**  
J.L. Kirk, J.N. Klironomos, H. Lee, J.T. Trevors  
*Plant-specific changes in microbial populations on roots affect degradation of petroleum hydrocarbons in contaminated soil.*
- 467 Quantifying the role of forest soil and bedrock in the acid neutralization of surface water in steep hillslopes**  
Y. Asano, T. Uchida  
*Bedrock plays a major role in neutralizing acid when overlying soils have been leached of base cations.*
- 481 Mercury burdens in Chinese mitten crabs (*Eriocheir sinensis*) in three tributaries of southern San Francisco Bay, California, USA**  
C.A. Hui, D. Rudnick, E. Williams  
*Hepatopancreas concentrations of mercury decline with crab size, which may have important consequences for bio-magnification in food webs.*

