


Volume 134 No. 3 April 2005 ISSN 0269-7491



ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning


ASSOCIATE EDITORS

K.E. Havens

S.V. Krupa

K.C. Jones

J. W. Erisman



ENVIRONMENTAL POLLUTION

CONTENTS—Continued from outside back cover

- 465 **Nitrogen content, ¹⁵N natural abundance and biomass of the two pleurocarpous mosses *Pleurozium schreberi* (Brid.) Mitt. and *Scleropodium purum* (Hedw.) Limpr. in relation to atmospheric nitrogen deposition**
A. Solga, J. Burkhardt, H.G. Zechmeister, J.-P. Frahm
Mosses are useful as monitors of nitrogen deposition.
- 475 **Relationships between reproductive performance and organochlorine contaminants in great black-backed gulls (*Larus marinus*)**
M. Helberg, J.O. Bustnes, K.E. Erikstad, K.O. Kristiansen, J.U. Skaare
Elevated blood concentrations of organochlorine contaminants correlate with poor reproductive performance in female great black-backed gulls.
- 485 **Biosensor-based diagnostics of contaminated groundwater: assessment and remediation strategy**
J. Bhattacharyya, D. Read, S. Amos, S. Dooley, K. Killham, G.I. Paton
Luminescent bacteria complement chemical analysis and support remediation technology.
- 493 **Metal partitioning and uptake in central Ontario forests**
S.A. Watmough, P.J. Dillon, E.N. Epova
Metal cycling by plants should be considered in risk assessment studies.
- 503 **Specific accumulation of 20 trace elements in great cormorants (*Phalacrocorax carbo*) from Japan**
D.-H. Nam, Y. Anan, T. Ikemoto, Y. Okabe, E.-Y. Kim, A. Subramanian, K. Saeki, S. Tanabe
Dietary sources seemed to be the main reason for variable results.
- 515 **Effects of arbuscular mycorrhizal fungi on the root uptake and translocation of radiocaesium**
H. Dupré de Boulois, B. Delvaux, S. Deçlerck
Radiocaesium root uptake and translocation is affected by an arbuscular mycorrhizal fungus.
- 525 **Degradation and leaching of the herbicides metolachlor and diuron: a case study in an area of Northern Italy**
A.B. Caracciolo, G. Giuliano, P. Grenni, L. Guzzella, F. Pozzoni, P. Bottoni, L. Fava, A. Crobe, M. Orrù, E. Funari
Pesticide mobility to vulnerable groundwaters in Italy is assessed and ranked.
- 535 **Practical considerations for addressing uncertainties in monitoring bulk deposition**
U. Dämmgen, J.W. Erisman, J.N. Cape, L. Grünhage, D. Fowler
Recommendations are given for sampling strategies and for making corrections and substitution of missing data.
- 549 **Spatial variation of mercury levels in nesting Bonelli's eagles from Southwest Portugal: effects of diet composition and prey contamination**
L. Palma, P. Beja, P.C. Tavares, L.R. Monteiro
The effects of diet composition and prey contamination added up to determine the spatial variation of Hg levels in breeding Bonelli's eagles.

CONTENTS

Volume 134 Number 3 2005

Research papers

- 363 Leaf litter processing and invertebrate assemblages along a pollution gradient in a Maine (USA) headwater stream**
T.S. Woodcock, A.D. Huryn
Variable pollution tolerance of community members provides stress resistance at the ecosystem level in streams.
- 377 No evidence for a critical salinity threshold for growth and reproduction in the freshwater snail *Physa acuta***
B.J. Kefford, D. Nugegoda
Responses of snails to increasing salinity were non-linear.
- 385 Amelioration of Indian urban air pollution phytotoxicity in *Beta vulgaris* L. by modifying NPK nutrients**
A. Singh, S.B. Agrawal, D. Rathore
*Air pollution caused adverse impact on growth and biomass accumulation of *Beta vulgaris* L. plants while higher fertility levels showed reduced yield losses.*
- 397 Bioaccumulation of PCBs in Arctic seabirds: influence of dietary exposure and congener biotransformation**
K. Borgå, H. Wolkers, J.U. Skaare, H. Hop, D.C.G. Muir, G.W. Gabrielsen
Contaminant patterns is linked to phylogeny and species-specific differences in enzyme systems and activity.
- 411 Performance of a constructed wetland treating intensive shrimp aquaculture wastewater under high hydraulic loading rate**
Y.-F. Lin, S.-R. Jing, D.-Y. Lee, Y.-F. Chang, Y.-M. Chen, K.-C. Shih
A constructed wetland was found to be technically and economically feasible for managing water quality of an intensive recirculating aquaculture system.
- 423 Tissue distributions and seasonal dynamics of the hepatotoxic microcystins-LR and -RR in a freshwater snail (*Bellamya aeruginosa*) from a large shallow, eutrophic lake of the subtropical China**
J. Chen, P. Xie, L. Guo, L. Zheng, L. Ni
Snails bioaccumulate microcystins in their hepatopancreas, gut and gonad, posing a risk to human consumers.
- 431 Aerobic biodegradation of organotin compounds in activated sludge batch reactors**
A.S. Stasinakis, N.S. Thomaidis, A. Nikolaou, A. Kantifes
The fate of organotins is assessed in activated sludge systems.
- 439 Ozone slows stomatal response to light and leaf wounding in a Mediterranean evergreen broadleaf, *Arbutus unedo***
E. Paoletti
In response to ozone exposure, stomata were slower in closing rather than in opening.
- 447 Transfer of selenium from prey to predators in a simulated terrestrial food chain**
W.A. Hopkins, B.P. Staub, J.A. Baionno, B.P. Jackson, L.G. Talent
Partitioning of selenium among tissues differs between male and female lizards.
- 457 Influence of pH on pesticide sorption by soil containing wheat residue-derived char**
G. Sheng, Y. Yang, M. Huang, K. Yang
Wheat char was effective for adsorption of pesticides in soil, with efficacy varying with pH and particular pesticides.