


Volume 135 No. 3 June 2005 ISSN 0269-7491



ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning


ASSOCIATE EDITORS

K.E. Havens

S.V. Krupa

K.C. Jones

J. W. Erisman



ENVIRONMENTAL POLLUTION

CONTENTS—Continued from outside back cover

445 Improved daily precipitation nitrate and ammonium concentration models for the Chesapeake Bay Watershed

J.W. Grimm, J.A. Lynch

A linear least-squares regression approach was used to develop daily precipitation nitrate and ammonium concentration models for the Chesapeake Bay Watershed.

Research Papers

457 Chemometrics methods for the investigation of methylmercury and total mercury contamination in mollusks samples collected from coastal sites along the Chinese Bohai Sea

W. Yawei, L. Liná, S. Jianbo, J. Guibin

Rapana venosa might be used as a potential biomonitor for Hg pollution in the Bohai Sea, China.

469 Effects of increased deposition of atmospheric nitrogen on an upland *Calluna* moor: N and P transformations

M.G. Pilkington, S.J.M. Caporn, J.A. Carroll, N. Cresswell, J.A. Lee, B.A. Emmett, D. Johnson

There are low rates of nitrogen loss from a moor ecosystem under both ambient and high atmospheric loading of nitrogen.

481 Risk to breeding success of waterbirds by contaminants in Hong Kong: evidence from trace elements in eggs

J.C.W. Lam, S. Tanabe, M.H.W. Lam, P.K.S. Lam

Analysis of trace element concentrations in eggs indicated that mercury and selenium might pose some risk to waterbirds in Hong Kong.

491 Equilibrium partitioning of ¹⁴C-benzo(a)pyrene and ¹⁴C-beñazolin between fractionated phases from an arable topsoil

J.-M. Séquaris, A. Lavorenti, P. Burauel

The partition coefficient (K_d) of chemicals was calculated according to two-phase or three-phase partitioning models.

501 The use of autecological and environmental parameters for establishing the status of lichen vegetation in a baseline study for a long-term monitoring survey

S. Gombert, J. Asta, M.R.D. Seaward

Ecological clustering which establishes the status of lichen vegetation can be reliably used as a baseline study to monitor environmental changes.

515 Soil remediation: humic acids as natural surfactants in the washings of highly contaminated soils

P. Conte, A. Agretto, R. Spaccini, A. Piccolo

Solutions of natural humic acids appear to be a better choice for washing highly polluted soils

CONTENTS

Volume 135 Number 3 2005

SPECIAL ISSUE SECTION: The National Atmospheric Deposition Program (25th Anniversary) and Ammonia Workshop

- 343 The National Atmospheric Deposition Program: 25 years of monitoring in support of science and policy: An ammonia workshop: the state of science and future needs**
J.A. Lynch, M. Kerchner
- 347 Spatial and temporal trends of precipitation chemistry in the United States, 1985–2002**
C.M.B. Lehmann, V.C. Bowersox, S.M. Larson
Data from the National Atmospheric Deposition Program National Trends Network indicate that significant changes in precipitation chemistry have occurred from 1985 to 2002.
- 363 Trends in atmospheric ammonium concentrations in relation to atmospheric sulfate and local agriculture**
V.R. Kelly, G.M. Lovett, K.C. Weathers, G.E. Likens
Relationships between NH_4^+ and SO_4^{2-} suggest that aerosol NH_4^+ may be more strongly linked to SO_2 emissions than to local agricultural NH_3 emissions.
- 371 Nonlinear regression and ARIMA models for precipitation chemistry in East Central Florida from 1978 to 1997**
D.M. Nickerson, B.C. Madsen
Autoregressive integrated moving average models of precipitation data are an improvement over nonlinear models for the prediction of precipitation chemistry composition.
- 381 Overview and assessment of techniques to measure ammonia emissions from animal houses: the case of the Netherlands**
J. Mosquera, G.J. Monteny, J.W. Erisman
There are no methods to measure ammonia effectively from outdoor stockyards.
- 389 Measuring ammonia emissions from land applied manure: an intercomparison of commonly used samplers and techniques**
T.H. Misselbrook, F.A. Nicholson, B.J. Chambers, R.A. Johnson
Three commonly used ammonia emission measurement techniques were compared for different manure types and gave CVs between 23 and 74%.
- 399 Managing ammonia emissions from livestock production in Europe**
J. Webb, H. Menzi, B.F. Pain, T.H. Misselbrook, U. Dämmgen, H. Hendriks, H. Döhler
Reducing NH_3 emissions following spreading of manures to land ranks highly because of the large abatement potential and relatively small cost.
- 407 Spatial and temporal variability of the overall error of National Atmospheric Deposition Program measurements determined by the USGS collocated-sampler program, water years 1989–2001**
G.A. Wetherbee, N.E. Latysh, J.D. Gordon
Spatial and temporal characteristics and confidence limits associated with the overall absolute error in NADP/NTN measurements are identified and quantified.
- 419 Deposition monitoring networks: what monitoring is required to give reasonable estimates of ammonia/ammonium?**
J.W. Erisman, A. Hensen, J. Mosquera, M. Sutton, D. Fowler
Current measurement methods allow monitoring of the deposition of ammonia and ammonium in a network consisting of core sites and low cost sites.
- 433 Atmospheric nitrogen inputs to the Delaware Inland Bays: the role of ammonia**
J.R. Scudlark, J.A. Jennings, M.J. Roadman, K.B. Savidge, W.J. Ullman

Local emissions from poultry production appear to significantly contribute to wet and dry atmospheric NH_x loading to the Delaware Inland Bays.