


Volume 139 No. 3 February 2006 ISSN 0269-7491



ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning


ASSOCIATE EDITORS

K.E. Havens

S.V. Krupa

K.C. Jones

J. W. Erisman



ENVIRONMENTAL POLLUTION

CONTENTS—Continued from outside back cover

498 Stabilization of mercury-containing wastes using sulfide

H. Piao, P.L. Bishop

Sulfide is an effective reagent for the stabilization of mercury when optimum stabilization pH and sulfide dosage are used.

507 Heavy metal speciation and phytotoxic effects of three representative sewage sludges for agricultural uses

I. Walter, F. Martínez, V. Cala

Different sewage sludge processing methods affected the distribution of metals and the seed germination test in different ways.

515 Tradescantia micronucleus test indicates genotoxic potential of traffic emissions in European cities

A. Klumpp, W. Ansel, G. Klumpp, V. Calatayud, J.P. Garrec, S. He, J. Peñuelas, A. Ribas, H. Ro-Poulsen, S. Rasmussen, M.J. Sanz, P. Vergne

The Tradescantia micronucleus test can be used to assess genotoxic potential at urban sites.

523 GC and GC-MS characterization of crude oil transformation in sediments and microbial mat samples after the 1991 oil spill in the Saudi Arabian Gulf coast

T. García de Oteyza, J.O. Grimalt

Cyanobacterial mats inhibit degradation of oil by reducing exposure to the atmosphere and seawater.

532 Lac Dufault sediment core trace metal distribution, bioavailability and toxicity to *Hyalella azteca*

M. Nowierski, D.G. Dixon, U. Borgmann

Toxicity and bioaccumulation tests with sediment cores provide more information on biological effects of metals than surface sediment tests.

541 Effect of copper on the adsorption of *p*-nitrophenol onto soils

Z.-g. Pei, X.-q. Shan, B. Wen, S. Zhang, L. Yan, S.U. Khan

*The adsorption of *p*-nitrophenol onto soils decreased in the presence of copper due to its competition for the adsorption sites.*

550 A microcosm test of adaptation and species specific responses to polluted sediments applicable to indigenous chironomids (Diptera)

S. Bahrndorff, J. Ward, V. Pettigrove, A.A. Hoffmann

A field test for adaptation applicable to indigenous chironomids suggests adaptation to metal contaminants in one species but not in other species.

561 Tissues and hair residues and histopathology in wild rats (*Rattus rattus* L.) and Algerian mice (*Mus spretus* Lataste) from an abandoned mine area (Southeast Portugal)

R. Pereira, M.L. Pereira, R. Ribeiro, F. Gonçalves

The bioaccumulation of As and Cd and signs of renal histopathological injury proved the value of Algerian mice as a bioindicator species in the risk assessment of contaminated sites.

CONTENTS

Volume 139 Number 3 2006

- 385 Comparative acute toxicity of organic pollutants and reference values for crustaceans. I. Branchiopoda, Copepoda and Ostracoda**
F. Sánchez-Bayo
A reference guide for the effects of chemicals on crustaceans.
- 421 Growth and photosynthetic responses of two pine species (*Pinus koraiensis* and *Pinus rigida*) in a polluted industrial region in Korea**
D.S. Choi, M. Kayama, H.O. Jin, C.H. Lee, T. Izuta, T. Koike
Pinus koraiensis seems to be more pollution tolerant than Pinus rigida.
- 433 Effect of bone char application on Pb bioavailability in a Pb-contaminated soil**
S.-B. Chen, Y.-G. Zhu, Y.-B. Ma, G. McKay
Bone char amendments show potential for remediation of Pb-contaminated soils.
- 440 Afforestation, seasalt episodes and acidification – A paired catchment study in western Norway**
T. Larssen, J. Holme
Seasalt episodes cause higher mobilization of toxic aluminum in sites afforested with spruce.
- 451 Avoidance of low doses of naphthalene by Collembola**
L. Boitaud, S. Salmon, C. Bourlette, J.-F. Ponge
PAH avoidance by soil springtails is species-specific and differs among populations of the same species.
- 455 The influence of single and multiple applications of pyrene on the evolution of pyrene catabolism in soil**
C.J.A. Macleod, K.T. Semple
Evolution of catabolic activity in soil is influenced by the concentration and number of applications of organic contaminants.
- 461 Records of atmospheric delivery of pyrolysis-derived pollutants in recent mountain lake sediments of the Julian Alps (NW Slovenia)**
G. Muri, S.G. Wakeham, N.L. Rose
Fluxes of black carbon, polycyclic aromatic hydrocarbons and spheroidal carbonaceous particles were determined in recent remote alpine lake sediments.
- 469 Genetic toxicity in surface water from Guaiba Hydrographic Region under the influence of industrial, urban and agricultural sewage in the *Drosophila* Wing-Spot Test**
V.S. do Amaral, M. Sinigaglia, M.L. Reguly, H.H.R. de Andrade
Drosophila Wing-Spot Test can be used for detection of environmental mutagenesis.
- 477 An interdisciplinary physical-chemical approach for characterization of arsenic in a calciner residue dump in Cornwall (UK)**
J.T. van Elteren, Z. Šlejkovec, I. Arčon, H.-J. Glass
Arsenic in calciner residue dumps is present as arsenate in crystalline ($Fe_2(As(AsO_4)_3)$) and amorphous or poorly-crystalline (bound to hematite and quartz) structures.
- 489 Determination of potential sources of PCBs and PBDEs in sediments of the Niagara River**
F. Samara, C.W. Tsai, D.S. Aga
Wastewater treatment plant discharges are a main source of PCBs and PBDEs to Niagara River sediments.