

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens K.C. Jones

S.V. Krupa

J.W. Erisman

ENVIRONMENTAL POLLUTION

CONTENTS-Continued from outside back cover

476 Tissue-specific accumulation and lactational transfer of polychlorinated biphenyls, chlorinated pesticides, and brominated flame retardants in hooded seals (*Cistophora cristata*) from the Gulf of St. Lawrence: Applications for monitoring

H. Wolkers, M.O. Hammill, B. van Bavel

Blood plasma contaminant concentrations differ from those in other tissues and might therefore be unsuitable for contaminant monitoring in hooded seals.

487 Presence of plutonium contamination in soils from Palomares (Spain)

M.C. Jiménez-Ramos, R. García-Tenorio, I. Vioque, G. Manjón, M. García-León

The remaining transuranic contamination in Palomares soils is mainly present in particulate form.

493 Linking biosensor responses to Cd, Cu and Zn partitioning in soils J.J.C. Dawson, C.D. Campbell, W. Towers, C.M. Cameron, G.I. Paton

Biosensors link biological hazard assessments of metals in soils with physico-chemical partitioning.

501 Using multivariate analyses and GIS to identify pollutants and their spatial patterns in urban soils in Galway, Ireland C. Zhang

Multivariate statistics and GIS are useful tools to identify pollutants in urban soils.

512 Characterization of contamination in and toxicities of a shipyard area in Hong Kong S.W. Chiu, K.M. Ho, S.S. Chan, O.M. So, K.H. Lai

Heavily contaminated shipyards result in high ecotoxicities but the environmental conditions handicap natural degradation.

521 Study of the trace metal ion influence on the turnover of soil organic matter in cultivated contaminated soils C. Dumat, K. Quenea, A. Bermond, S. Toinen, M.F. Benedetti

Measure of δ^{13} C and total metal concentrations in size fractions of contaminated soils suggests an influence of metals on the soil organic matter dynamic.

530 Field evaluation of in situ remediation of a heavy metal contaminated soil using lime and red-mud C.W. Gray, S.J. Dunham, P.G. Dennis, F.J. Zhao, S.P. McGrath

Red mud was effective in immobilising heavy metals in soil.

Physiological and biochemical stress responses in grassland species are influenced by both early-season ozone exposure and interspecific competition

F. Scebba, F. Canaccini, A. Castagna, J. Bender, H.-J. Weigel, A. Ranieri

The competition between species modulates the ozone effect in a short-term.

Environmental contamination of mercury from Hg-mining areas in Wuchuan, northeastern Guizhou, China G. Qiu, X. Feng, S. Wang, L. Shang

Mercury mining activities in Wuchun, Guizhou, China have resulted in seriously mercury contamination to the local environment.

559 Effects of eutrophication on vitellogenin gene expression in male fathead minnows (Pimephales promelas) exposed to 17α-ethynylestradiol in field mesocosms

D.A. Gordon, G.P. Toth, D.W. Graham, J.M. Lazorchak, T.V. Reddy, C.W. Knapp, F. deNoyelles Jr., S. Campbell, D.L. Lattier

Nutrient condition in surface waters strongly influences vitellogenin gene expression in male fathead minnows.

Geostatistical analysis and kriging of Hexachlorocyclohexane residues in topsoil from Tianjin, China B.G. Li, J. Cao, W.X. Liu, W.R. Shen, X.J. Wang, S. Tao

Geostatistical analysis and kriging were applied to HCH concentrations in topsoil of Tianjin, China for mapping purposes.

ENVIRONMENTAL POLLUTION

www.elsevier.com/locate/envpol

CONTENTS

Volume 142 Number 3 2006

383 Reproductive transfer of organochlorines in viviparous surfperch, Ditrema temmincki M. Oka, T. Arai, Y. Shibata, N. Miyazaki

The reproductive transfer of POPs in matrotrophic viviparous surfperch was determined for the first time.

388 Atmospheric polycyclic aromatic hydrocarbons (PAHs) in Asia: A review from 1999 to 2004 K.-F. Chang, G.-C. Fang, J.-C. Chen, Y.-S. Wu

This work summarizes the characteristics, sources and distribution, and the healthy impacts of atmospheric PAH species in Asia.

397 Selenium and other trace metals in fish inhabiting a fly ash stream: Implications for regulatory tissue thresholds R.J. Reash, T.W. Lohner, K.V. Wood

Bluegill sunfish inhabiting a coal fly ash receiving stream had elevated selenium levels in whole body and gonad tissue (9–10 times higher than reference fish), and antagonistic metal interactions may be one of several mechanisms allowing long-term persistence of the population.

409 Geochemical and Pb isotopic evidence for sources and dispersal of metal contamination in stream sediments from the mining and smelting district of Příbram, Czech Republic
V. Ettler, M. Mihaljevič, O. Šebek, M. Molek, T. Grygar, J. Zeman

Pb isotopes properly complete traditional investigations of metal sources and dispersal in contaminated stream sediments.

418 Mercury accumulation in surface sediments of salt marshes of the Bay of Fundy G.A. Hung, G.L. Chmura

Short-term storage of mercury was measured in salt marsh sediments of the Bay of Fundy.

432 Eco-toxicological studies of diesel and biodiesel fuels in aerated soil A. Lapinskienė; P. Martinkus, V. Rėbždaitė

The comparison of diesel and biodiesel fuels' eco-toxicological parameters in non-adapted aerated soil is relevant when considering the effects of these substances on the environment in cases of accidental spills.

438 Non-destructive pollution exposure assessment in the European hedgehog (*Erinaceus europaeus*): II. Hair and spines as indicators of endogenous metal and As concentrations
H. D'Havé, J. Scheirs, V.K. Mubiana, R. Verhagen, R. Blust, W. De Coen

Hedgehog hair and spines are promising non-destructive biomonitoring tools of terrestrial metal pollution.

449 Cosorption study of organic pollutants and dissolved organic matter in a soil F. Flores-Céspedes, M. Fernández-Pérez, M. Villafranca-Sánchez, E. González-Pradas

Cosorption of organic pollutants and DOM.

457 High incidence of deformity in aquatic turtles in the John Heinz National Wildlife Refuge B. Bell, J.R. Spotila, J. Congdon

This paper presents findings on the prevalence of developmental abnormalities in turtles at a national wildlife refuge that have direct relevance to studies on the effects of contamination on development and morphology of vertebrates.

466 Polycyclic aromatic hydrocarbons affect survival and development of common snapping turtle (Chelydra serpentina) embryos and hatchlings

R.J. Van Meter, J.R. Spotila, H.W. Avery

Exposure to polycyclic aromatic hydrocarbons on the egg reduces survival of snapping turtle embryos and causes developmental abnormalities.