

ENVIRONMENTAL POLLUTION


EDITOR-IN-CHIEF

W.J. Manning

ASSOCIATE EDITORS

K.E. Havens S.V. Krupa

K.C. Jones J.W. Erisman



ENVIRONMENTAL POLLUTION

www.elsevier.com/locate/envpol

CONTENTS

Volume 143 Number 1 2006

Letter to the editor

In defence of plants as biomonitors of soil quality P. Madejón, T. Marañón, J.M. Murillo, B. Robinson

The use of plants as biomonitors of soil quality has important advantages, particularly on a large scale.

Regular papers

4 Cadmium and lead concentrations in Skrjabinotaenia lobata (Cestoda: Catenotaeniidae) and in its host, Apodemus sylvaticus (Rodentia: Muridae) in the urban dumping site of Garraf (Spain)
J. Torres, J. Peig, C. Eira, M. Borrás

The parasitological model S. lobata/A. sylvaticus presents suitable features to be used as a bioindicator of lead pollution in terrestrial habitats.

9 Effects of pH, organic acids, and inorganic ions on lead desorption from soils J.Y. Yang, X.E. Yang, Z.L. He, T.Q. Li, J.L. Shentu, P.J. Stoffella

The adsorption-desorption process is a basic and important reaction in soils controlling Pb2+ mobility and bioavailability.

A comparison of Cu, Pb, As, Cd, Zn, Fe, Ni and Mn determined by acid extraction/ICP-OES and ex situ field portable X-ray fluorescence analyses
C. Kilbride, J. Poole, T.R. Hutchings

Portable X-ray fluorescence instruments can provide an accurate, rapid and cost-effective means of assessing soil contamination and for monitoring the efficacy of remediation strategies.

Distribution of sorbed phenanthrene and pyrene in different humic fractions of soils and importance of humin B. Pan, B.S. Xing, W.X. Liu, S. Tao, X.M. Lin, X.M. Zhang, Y.X. Zhang, Y. Xiao, H.C. Dai, H.S. Yuan

Humic fractionation after sorption experiment revealed that humin fraction is the main region for slow and nonlinear sorption of phenanthrene and pyrene.

34 Assessment of acid leachable trace metals in sediment cores from River Uppanar, Cuddalore, Southeast coast of India T. Ayyamperumal, M.P. Jonathan, S. Srinivasalu, J.S. Armstrong-Altrin, V. Ram-Mohan

Both natural and anthropogenic factors are affecting metals in sediments.

46 Source determination of highly chlorinated biphenyl isomers in pine needles – Comparison to several PCB preparations B. Wyrzykowska, I. Bochentin, N. Hanari, A. Orlikowska, J. Falandysz, H. Yuichi, N. Yamashita

Pine needles were suitable as passive samplers for PCBs in air.

60 A predictive model for copper partitioning to suspended particulate matter in river waters Y. Lu, H.E. Allen

Cu partitioning is modeled in natural water systems with different pH, alkalinity, and dissolved and particulate organic carbon, suspended solids, and copper concentrations.

73 EPR monitoring of the bioavailability of an organic xenobiotic (4-hydroxy-TEMPO) in model clay suspensions and pastes A. Dumestre, M. Spagnuolo, R. Bladon, J. Berthelin, P. Baveye


Spectroscopic data demonstrate that the bioavailability of an organic compound in clay suspensions and pastes strongly depends on conditions that affect bacterial metabolism.

81 Biomonitoring using the lichen *Hypogymnia physodes* and bark samples near Zlatna, Romania immediately following closure of a copper ore-processing plant
A.-M. Rusu, G.C. Jones, P.D.J. Chimonides, O.W. Purvis

Lichen transplants and bark samples were useful as indicators of short- and long-term particulate and gaseous deposition.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; Index Medicus/MEDLINE/PubMed; ISI GeoSciTech; Science Citation Index; SciSearch)



0269-7491(200609)143:1;1-S

