

Volume 143 No. 3 October 2006 ISSN 0269-7491

ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

W.J. Manning


ASSOCIATE EDITORS

K.E. Havens

S.V. Krupa

K.C. Jones

J.W. Erisman



CONTENTS

Volume 143 Number 3 2006

- 377 Detecting changes in insect herbivore communities along a pollution gradient**
M.E. Jones, T.D. Paine
Differences in insect herbivore communities were associated with an ambient air pollution gradient in the mixed conifer forest outside the Los Angeles area.
- 388 Abiotic ecotypes in south-central Spanish rivers: Reference conditions and pollution**
J.L. Moreno, C. Navarro, J. De las Heras
Agriculture by means of nutrient surpluses and water diversion for irrigation, along with poor sewage treatment of urban wastes are the main environmental problems in Spanish rivers.
- 397 A remediation strategy based on active phytoremediation followed by natural attenuation in a soil contaminated by pyrite waste**
R. Clemente, C. Almela, M.P. Bernal
The addition of lime and organic amendments decreased heavy metal solubility and promoted the natural attenuation of a recently-contaminated soil.
- 407 Fixation of metals in soil constituents and potential remobilization by hyperaccumulating and non-hyperaccumulating plants: Results from an isotopic dilution study**
D. Hammer, C. Keller, M.J. McLaughlin, R.E. Hamon
Thlaspi caerulescens takes up more Cd and Zn than Brassica napus but from the same pools.
- 416 The use of soil survey data to determine the magnitude and extent of historic metal deposition related to atmospheric smelter emissions across Humberside, UK**
B.G. Rawlins, R.M. Lark, R. Webster, K.E. O'Donnell
Soil survey data are used to estimate the deposition of metals to land surrounding a former smelter.
- 427 Seasonal profiles of leaf ascorbic acid content and redox state in ozone-sensitive wildflowers**
K.O. Burkey, H.S. Neufeld, L. Souza, A.H. Chappelka, A.W. Davison
Wildflower species exhibit differences in ascorbic acid content and redox status that affect ozone sensitivity.
- 435 Distribution and relationships between selected chemical elements in green alga *Enteromorpha* sp. from the southern Baltic**
R. Żbikowski, P. Szefer, A. Latała
Enteromorpha sp. can be used as efficient biomonitor for chemical elements in coastal areas of the Baltic Sea.
- 449 Cry1Ab protein from Bt transgenic rice does not residue in rhizosphere soil**
H. Wang, Q. Ye, W. Wang, L. Wu, W. Wu
Cry1Ab protein was not detected in the rhizosphere soil of field-grown Bt transgenic rice.
- 456 Contamination in sediments, bivalves and sponges of McMurdo Sound, Antarctica**
A. Negri, K. Burns, S. Boyle, D. Brinkman, N. Webster
Analyses of Antarctic marine sediments, bivalves and sponges revealed strong PAH, PCB and trace metal gradients in McMurdo Sound.
- 468 Modelling nitrogen saturation and carbon accumulation in heathland soils under elevated nitrogen deposition**
C.D. Evans, S.J.M. Caporn, J.A. Carroll, M.G. Pilkington, D.B. Wilson, N. Ray, N. Cresswell
Enhanced carbon sequestration may slow the rate of nitrogen saturation in heathlands.
- 479 Changes in elemental composition and mass of atmospheric aerosol pollution between 1996 and 2002 in a Central European city**
I. Salma, W. Maenhaut
Non-exhaust traffic emission particles and coarse-mode particles are increasing in Budapest, Hungary.

continued on inside back cover

(Abstracted/indexed in: AGRICOLA database; Air Pollution Control Association Journal; Biological and Agricultural Index; CAB ABSTRACTS database; Elsevier BIOBASE/Current Awareness in Biological Sciences; Cambridge Scientific Abstracts; Chemical Abstracts; Current Contents/Agriculture, Biology & Environmental Sciences; Environment Abstracts; Environmental Periodicals Bibliography; Energy Information Abstracts; EMBASE/Excerpta Medica; Geo Abstracts; GEOBASE; Index Medicus/MEDLINE/PubMed; ISI GeoSciTech; Science Citation Index; SciSearch)



0269-7491(200610)143:3;1-W



ELSEVIER

02022