


60218

516

MONOCYTE-REACTIVE IgG ALLOANTIBODIES
IN Rh D-HAEMOLYTIC DISEASE
OF THE NEWBORN



Marion Dooren

CONTENTS

Chapter I		
Introduction		1
Chapter II		
Predictive value of a monocyte-driven cytotoxicity assay for the severity of Rh D-haemolytic disease of the newborn		7
Chapter III		
Detection of human monocyte-reactive alloantibodies by flow cytometry after selective downmodulation of the Fc receptor I <i>Blood</i> 1991;78:2150-2156		19
Chapter IV		
Protection against immune haemolytic disease of newborn infants by maternal monocyte-reactive IgG alloantibodies (anti-HLA-DR). <i>Lancet</i> 1992;339:1067-1070		35
Chapter V		
Evidence for the protective effect of maternal FcR-blocking IgG alloantibodies (HLA DR) in Rh D-haemolytic disease of the newborn		45
Chapter VI		
Protection against Rh D-haemolytic disease of the newborn by a diminished transport of maternal IgG to the fetus.		55
Chapter VII		
Maternal monocyte-reactive IgG alloantibodies in Rh D-haemolytic disease of the newborn. I. Their specificity and frequency of occurrence.		61

Chapter VIII

Maternal monocyte-reactive IgG alloantibodies in Rh D-haemolytic disease of the newborn.

II. Their mechanism of action

73

Chapter IX

Adult respiratory distress syndrome after administration of an experimental intravenous gammaglobulin concentrate with a high titre of monocyte-reactive IgG antibodies

85

Chapter X

No beneficial effect of intravenous gammaglobulin in combination with intravascular fetal transfusions in severe Rh D-haemolytic disease

101

Summary

111

Samenvatting

115