


Volume 133 No. 2 January 2005 ISSN 0269-7491



# ENVIRONMENTAL POLLUTION

EDITOR-IN-CHIEF

**W.J. Manning**


ASSOCIATE EDITORS

**K.E. Havens**

**S.V. Krupa**

**K.C. Jones**

**J. W. Erisman**



# ENVIRONMENTAL POLLUTION

## CONTENTS—Continued from outside back cover

- 283 **Polycyclic aromatic hydrocarbons storage by *Fusarium solani* in intracellular lipid vesicles**  
A. Verdin, A. Lounès-Hadj Sahraoui, R. Newsam, G. Robinson, R. Durand  
*Fungi can store PAHs intracellularly in lipid vesicles independently of their PAH degradation abilities.*
- 293 ***Pinus nigra* and *Pinus pinaster* needles as passive samplers of polycyclic aromatic hydrocarbons**  
M.T. Piccardo, M. Pala, B. Bonaccorso, A. Stella, A. Redaelli, G. Paola, F. Valerio  
*The suitability of the pine needles as passive air samplers for persistent trace organics is demonstrated.*
- 303 **UV-absorbing compounds in subarctic herbarium bryophytes**  
S. Huttunen, N.M. Lappalainen, J. Turunen  
*UV-B-absorbing compounds in subarctic herbarium bryophytes indicate the radiation climate of the collecting site and time.*
- 315 **Environmental specimen bank samples of *Pleurozium schreberi* and *Hylocomium splendens* as indicators of the radiation environment at the surface**  
S. Huttunen, T. Taipale, N.M. Lappalainen, E. Kubin, K. Lakkala, J. Kaurola  
*Acidified methanol-soluble UV-B-absorbing compounds of environmental specimen bank samples indicated the surface radiation environment of the collection site and season.*
- 327 **Uptake of radionuclides by vegetation at a High Arctic location**  
M. Dowdall, J.P. Gwynn, C. Moran, J. O'Dea, C. Davids, B. Lind  
*Uptake of the anthropogenic radionuclide <sup>137</sup>Cs is highest for moss species.*
- 333 **The significance of water ionic strength on aluminium toxicity in brown trout (*Salmo trutta* L.)**  
N.E.W. Alstad, B.M. Kjelsberg, L.A. Vøllestad, E. Lydersen, A.B.S. Poléo  
*Ionic strength has a slight ameliorating effect on Al toxicity in brown trout.*
- 343 **Age dependence of the accumulation of organochlorine pollutants in brown trout (*Salmo trutta*) from a remote high mountain lake (Redó, Pyrenees)**  
I. Vives, J.O. Grimalt, M. Ventura, J. Catalan, B.O. Rosseland  
*Trout in high mountain lakes display age-dependent accumulation of certain organochlorine pollutants.*
- 351 **Modelling the transport of suspended particulate matter by the Rhone River plume (France). Implications for pollutant dispersion**  
R. Periañez  
*A model has been developed to simulate transport of suspended particulate matter in the Rhone River Plume.*
- 365 **A comparative study of cadmium phytoextraction by accumulator and weed species**  
M. Ghosh, S.P. Singh  
**Ipomoea carnea* was more effective in extracting cadmium than was *Brassica juncea*.*
- 373 **Heavy metal accumulation, heat shock protein expression and cytogenetic changes in *Tetrix tenuicornis* (L.) (Tetrigidae, Orthoptera) from polluted areas**  
E. Warchałowska-Śliwa, M. Niklińska, A. Görlich, P. Michailova, E. Pyza  
*Metals accumulation caused genotoxicity in insects.*
- 383 **Study of PAH emission from the solid fuels combustion in residential furnaces**  
S.V. Kakareka, T.I. Kukharchyk, V.S. Khomich  
*Emission factors are reported for PAHs from residential solid fuel combustion.*

**CONTENTS**

**Volume 133 Number 2 2005**



**183 Environmental applications for biosurfactants**

C.N. Mulligan

*More information is needed to be able to predict and model the behaviour of biosurfactants.*

**199 Establishing principal soil quality parameters influencing earthworms in urban soils using bioassays**

P.K. Hankard, J.G. Bundy, D.J. Spurgeon, J.M. Weeks, J. Wright, C. Weinberg, C. Svendsen

*Bioassays must be applied for the risk assessment complexly-polluted sites to complement chemical analysis of soils.*

**213 The use of the barbell cluster ANOVA design for the assessment of environmental pollution: a case study, Wigierski National Park, NE Poland**

Z.M. Migaszewski, A. Galuszka, P. Paslawski

*The barbell cluster ANOVA design allowed the number of samples collected to be reduced to a minimum.*

**225 Recycling EDTA solutions used to remediate metal-polluted soils**

K.R. Zeng, S. Sauvé, H.E. Allen, W.H. Hendershot

*EDTA could be recycled seven times over a 14-day period, enhancing removal of metals from soils.*

**233 Zn, Cd and Pb accumulation and arbuscular mycorrhizal colonisation of pennycress *Thlaspi praecox* Wulf. (Brassicaceae) from the vicinity of a lead mine and smelter in Slovenia**

K. Vogel-Mikuš, D. Drobne, M. Regvar

*Thlaspi praecox Wulf. (Brassicaceae) is a newly discovered Cd, Zn and Pb hyperaccumulator able to form symbiosis with arbuscular mycorrhizal fungi.*

**243 Effects of golf course construction and operation on water chemistry of headwater streams on the Precambrian Shield**

J.G. Winter, P.J. Dillon

*Golf course construction and operation had a significant impact on alkalinity, nitrogen and base cation concentrations of streams.*

**255 Priority volatile organic compounds in surface waters of the southern North Sea**

T. Huybrechts, J. Dewulf, H. Van Langenhove

*Volatile aromatics are a major group of volatile organic compounds in the North Sea, and are attributed to discharges from shipping and oil related activities.*

**265 Influence of temperature and salinity on heavy metal uptake by submersed plants**

Å. Fritioff, L. Kautsky, M. Greger

*Metal concentrations increase with increasing temperature and decreasing salinity in two aquatic plants.*

**275 Effects of some environmental parameters on catalase activity measured in the mussel (*Mytilus galloprovincialis*) exposed to lindane**

A. Khessiba, M. Roméo, P. Aïssa

*Oxidative stress in mussels exposed to lindane was also influenced by a number of abiotic parameters.*